

JEE MAIN + ADVANCED

MATHEMATICS

TOPIC NAME

CONTINUITY

&

DIFFERENTIABILITY

(PRACTICE SHEET)

LEVEL-1

Question based on

Continuity of a function at a point

- Q.1** Function $f(x) = \begin{cases} 1+x, & \text{when } x < 2 \\ 5-x, & \text{when } x > 2 \end{cases}$; $x = 2$ is continuous at $x = 2$, if $f(2)$ equals -
 (A) 0 (B) 1 (C) 2 (D) 3
- Q.2** If $f(x) = \begin{cases} x \cos \frac{1}{x}, & x \neq 0 \\ k, & x = 0 \end{cases}$ is continuous at $x = 0$, then
 (A) $k > 0$ (B) $k < 0$
 (C) $k = 0$ (D) $k \geq 0$
- Q.3** If function $f(x) = \begin{cases} x^2 + 2, & x > 1 \\ 2x + 1, & x = 1 \end{cases}$ is continuous at $x = 1$, then value of $f(x)$ for $x < 1$ is -
 (A) 3 (B) $1 - 2x$
 (C) $1 - 4x$ (D) None of these
- Q.4** Which of the following function is continuous at $x = 0$ -
 (A) $f(x) = \begin{cases} \sin \frac{2x}{x}, & x \neq 0 \\ 1, & x = 0 \end{cases}$
 (B) $f(x) = \begin{cases} (1+x)^{1/x}, & x \neq 0 \\ 1, & x = 0 \end{cases}$
 (C) $f(x) = \begin{cases} e^{-1/x}, & x \neq 0 \\ 1, & x = 0 \end{cases}$
 (D) None of these
- Q.5** If $f(x) = \begin{cases} 6 \times 5^x, & x \leq 0 \\ 2a + x, & x > 0 \end{cases}$ is continuous at $x = 0$, then the value of a is -
 (A) 1 (B) 2
 (C) 3 (D) None of these
- Q.6** If $f(x) = \begin{cases} \frac{x^2 - (a+2)x + a}{x-2}, & x \neq 2 \\ 2, & x = 2 \end{cases}$ is continuous at $x = 2$, then a is equal to-
 (A) 0 (B) 1 (C) -1 (D) 2
- Q.7** If $f(x) = \begin{cases} \frac{\sin^{-1} ax}{x}, & x \neq 0 \\ k, & x = 0 \end{cases}$ is continuous at $x = 0$, then k is equal to-
 (A) 0 (B) 1
 (C) a (D) None of these
- Q.8** What is the value of $(\cos x)^{1/x}$ at $x = 0$ so that it becomes continuous at $x = 0$ -
 (A) 0 (B) 1 (C) -1 (D) e
- Q.9** If $f(x) = \begin{cases} \frac{k \cos x}{\pi - 2x}, & x \neq \pi/2 \\ 1, & x = \pi/2 \end{cases}$ is a continuous function at $x = \pi/2$, then the value of k is-
 (A) -1 (B) 1 (C) -2 (D) 2
- Q.10** If function $f(x) = \frac{x^3 - a^3}{x - a}$, is continuous at $x = a$, then the value of $f(a)$ is -
 (A) $2a$ (B) $2a^2$ (C) $3a$ (D) $3a^2$
- Q.11** If $f(x) = \begin{cases} \sin \frac{1}{x}, & x \neq 0 \\ k, & x = 0 \end{cases}$ is continuous at $x = 0$, then k is equal to -
 (A) 8 (B) 1
 (C) -1 (D) None of these
- Q.12** Function $f(x) = \left(1 + \frac{x}{a}\right)^{1/x}$ is continuous at $x = 0$ if $f(0)$ equals-
 (A) e^a (B) e^{-a}
 (C) 0 (D) $e^{1/a}$

Q.13 If $f(x) = \frac{1 - \cos 7(x - \pi)}{x - \pi}$, ($x \neq \pi$) is continuous at $x = \pi$, then $f(\pi)$ equals -
 (A) 0 (B) 1 (C) -1 (D) 7

Q.14 If $f(x) = \begin{cases} \frac{\tan x}{\sin x}, & x \neq 0 \\ 1, & x = 0 \end{cases}$, then $f(x)$ is -
 (A) continuous everywhere
 (B) continuous nowhere
 (C) continuous at $x = 0$
 (D) continuous only at $x = 0$

Q.15 If $f(x) = \frac{2x + \tan x}{x}$ is continuous at $x = 0$, then $f(0)$ equals -
 (A) 0 (B) 1 (C) 2 (D) 3

Q.16 If $f(x) = \frac{\sqrt{1+x} - \sqrt[3]{1+x}}{x}$, ($x \neq 0$) is continuous at $x = 0$, then the value of $f(0)$ is -
 (A) 1/6 (B) 1/4 (C) 2 (D) 1/3

Q.17 If $f(x) = \begin{cases} ax^2 - b & \text{when } 0 \leq x < 1 \\ 2 & \text{when } x = 1 \\ x + 1 & \text{when } 1 < x \leq 2 \end{cases}$ is continuous at $x = 1$, then the most suitable values of a, b are -
 (A) $a = 2, b = 0$ (B) $a = 1, b = -1$
 (C) $a = 4, b = 2$ (D) All the above

Q.18 If $f(x) = \begin{cases} |x|, & \text{when } x < 0 \\ x, & \text{when } 0 \leq x < 1 \\ 1, & \text{when } x > 1 \end{cases}$ then f is -
 (A) continuous for every real number x
 (B) discontinuous at $x = 0$
 (C) discontinuous at $x = 1$
 (D) discontinuous at $x = 0$ and $x = 1$

Q.19 If $f(x) = \begin{cases} \sin(1/x), & x \neq 0 \\ 0, & x = 0 \end{cases}$, then it is discontinuous at -
 (A) $x = 0$ (B) All points
 (C) No point (D) None of these

Q.20 Function $f(x) = x - |x|$ is -
 (A) discontinuous at $x = 0$
 (B) discontinuous at $x = 1$
 (C) continuous at all points
 (D) discontinuous at all points

Q.21 Function $f(x) = \tan x$, is discontinuous at -
 (A) $x = 0$ (B) $x = \pi/2$
 (C) $x = \pi$ (D) $x = -\pi$

Q.22 Function $f(x) = [x]$ is discontinuous at -
 (A) every real number
 (B) every natural number
 (C) every integer
 (D) No where

Q.23 Function $f(x) = 3x^2 - x$ is -
 (A) discontinuous at $x = 1$
 (B) discontinuous at $x = 0$
 (C) continuous only at $x = 0$
 (D) continuous at $x = 0$

Q.24 If $f(x) = \begin{cases} x^2, & \text{when } x \leq 0 \\ 1, & \text{when } 0 < x < 1 \\ 1/x, & \text{when } x \geq 1 \end{cases}$, then $f(x)$ is -
 (A) continuous at $x = 0$ but not at $x = 1$
 (B) continuous at $x = 1$ but not at $x = 0$
 (C) continuous at $x = 0$ and $x = 1$
 (D) discontinuous at $x = 0$ and $x = 1$

Q.25 Function $f(x) = \begin{cases} -1, & x \in \mathbb{Q} \\ 1, & x \notin \mathbb{Q} \end{cases}$ is -
 (A) continuous at $x = 0$
 (B) continuous at $x = 1$
 (C) every where continuous
 (D) every where discontinuous

Q.26 If $f(x) = \begin{cases} -x^2, & x \leq 0 \\ 5x - 4, & 0 < x \leq 1 \\ 4x^2 - 3x, & 1 < x < 2 \\ 3x + 4, & x \geq 2 \end{cases}$, then $f(x)$ is -
 (A) continuous at $x = 0$ but not at $x = 1$
 (B) continuous at $x = 2$ but not at $x = 0$
 (C) continuous at $x = 0, 1, 2$
 (D) discontinuous at $x = 0, 1, 2$

Q.27 Function $f(x) = \begin{cases} \frac{x^2 - 4x + 3}{x^2 - 1}, & x \neq 1 \\ 2, & x = 1 \end{cases}$ is-

- (A) continuous at $x = 1$
- (B) continuous at $x = -1$
- (C) continuous at $x = 1$ and $x = -1$
- (D) discontinuous at $x = 1$ and $x = -1$

Q.28 Let $f(x) = 3 - |\sin x|$, then $f(x)$ is-

- (A) Everywhere continuous
- (B) Everywhere discontinuous
- (C) Continuous only at $x = 0$
- (D) Discontinuous only at $x = 0$

Q.29 The function $f(x) = \begin{cases} x-1, & x < 2 \\ 2x-3, & x \geq 2 \end{cases}$ is a

- continuous function for-
- (A) all real values of x
 - (B) only $x = 2$
 - (C) all real values of $x \neq 2$
 - (D) only all integral values of x

Q.30 If $f(x) = \begin{cases} x \sin x, & 0 < x \leq \pi/2 \\ \frac{\pi}{2} \sin(\pi + x), & \frac{\pi}{2} < x < \pi \end{cases}$, then -

- (A) $f(x)$ is discontinuous at $x = \pi/2$
- (B) $f(x)$ is continuous at $x = \pi/2$
- (C) $f(x)$ is continuous at $x = 0$
- (D) None of these

Q.31 The value of k so that

$$f(x) = \begin{cases} k(2x - x^2) & \text{when } x < 0 \\ \cos x, & \text{when } x \geq 0 \end{cases}$$

continuous at $x = 0$ is-

- (A) 1
- (B) 2
- (C) 4
- (D) None of these

Q.32 If $f(x) = \frac{(a+x)^2 \sin(a+x) - a^2 \sin a}{x}$, $x \neq 0$;

then the value of $f(0)$ so that f is continuous at $x = 0$ is-

- (A) $a^2 \cos a + a \sin a$
- (B) $a^2 \cos a + 2a \sin a$
- (C) $2a^2 \cos a + a \sin a$
- (D) None of these

Q.33 Let $f(x) = |x| + |x-1|$, then-

- (A) $f(x)$ is continuous at $x = 0$ and $x = 1$
- (B) $f(x)$ is continuous at $x = 0$ but not at $x = 1$
- (C) $f(x)$ is continuous at $x = 1$ but not at $x = 0$
- (D) None of these

Q.34 Consider the following statements:

- I. A function f is continuous at a point $x_0 \in \text{Dom}(f)$ if $\lim_{x \rightarrow x_0} f(x) = f(x_0)$.
- II. f is continuous in $[a, b]$ if f is continuous in (a, b) and $f(a) = f(b)$.
- III. A constant function is continuous in an interval.

Out of these correct statements are

- (A) I and II
- (B) II and III
- (C) I and III
- (D) All the above

Q.35 If $f(x) = \begin{cases} x+2, & \text{when } x < 1 \\ 4x-1, & \text{when } 1 \leq x \leq 3 \\ x^2+5, & \text{when } x > 3 \end{cases}$, then correct

statement is-

- (A) $\lim_{x \rightarrow 1} f(x) = \lim_{x \rightarrow 3} f(x)$
- (B) $f(x)$ is continuous at $x = 3$
- (C) $f(x)$ is continuous at $x = 1$
- (D) $f(x)$ is continuous at $x = 1$ and 3

Q.36 Let $f(x)$ and $\phi(x)$ be defined by $f(x) = [x]$ and

$$\phi(x) = \begin{cases} 0, & x \in I \\ x^2, & x \in \mathbb{R} - I \end{cases} \quad [.] = \text{G.I.F.}$$

- (A) $\lim_{x \rightarrow 1} \phi(x)$ exist but ϕ is not continuous at $x = 1$
- (B) $\lim_{x \rightarrow 1} f(x)$ does not exist and f is continuous at $x = 1$
- (C) ϕ is continuous for all x
- (D) None of these

Q.37 $f(x) = \begin{cases} \frac{x-4}{|x-4|} + a, & x < 4 \\ a+b, & x = 4 \\ \frac{x-4}{|x-4|} + b, & x > 4 \end{cases}$ is continuous at

$x = 4$, if-

- (A) $a = 0, b = 0$
- (B) $a = 1, b = 1$
- (C) $a = 1, b = -1$
- (D) $a = -1, b = 1$

Q.38 The function $f(x) = \frac{\cos x - \sin x}{\cos 2x}$ is continuous

everywhere then $f(\pi/4) =$

- (A) 1 (B) -1
(C) $\sqrt{2}$ (D) $1/\sqrt{2}$

Q.39 If $f(x) = \frac{\tan\left(\frac{\pi}{4} - x\right)}{\cot 2x}$, $x \neq \pi/4$ is every where

continuous, then $f(\pi/4)$ equals-

- (A) 0 (B) 1 (C) -1 (D) 1/2

Question based on

Continuity from left and right

Q.40 If $f(x) = \begin{cases} \frac{x}{e^{1/x} + 1}, & x \neq 0 \\ 0, & x = 0 \end{cases}$, then -

- (A) $\lim_{x \rightarrow 0^+} f(x) = 1$
(B) $\lim_{x \rightarrow 0^-} f(x) = 1$
(C) $f(x)$ is continuous at $x = 0$
(D) None of these

Q.41 If $f(x) = [x]$, where $[x] =$ greatest integer $\leq x$, then at $x = 1$, f is-

- (A) continuous (B) left continuous
(C) right continuous (D) None of these

Question based on

Continuity of a function in an interval

Q.42 If $f(x) = \begin{cases} \frac{\sqrt{1+px} - \sqrt{1-px}}{x}, & -1 \leq x < 0 \\ \frac{2x+1}{x-2}, & 0 \leq x \leq 1 \end{cases}$ is

continuous in the interval $[-1, 1]$ then p equals -

- (A) -1 (B) 1
(C) 1/2 (D) -1/2

Q.43 If $f(x) = \begin{cases} \frac{x^2}{a}, & 0 \leq x < 1 \\ a, & 1 \leq x < \sqrt{2} \\ \frac{(2b^2 - 4b)}{x^2}, & \sqrt{2} \leq x < \infty \end{cases}$ is

continuous in the interval $[0, \infty)$, then values of a and b are respectively -

- (A) 1, -1 (B) -1, $1 + \sqrt{2}$
(C) -1, 1 (D) None of these

Q.44 Which of the following function is not continuous in the interval $(0, \pi)$

- (A) $x \sin \frac{1}{x}$
(B) $\begin{cases} 1, & 0 < x \leq \frac{3\pi}{4} \\ 2 \sin\left(\frac{2x}{9}\right), & \frac{3\pi}{4} < x < \pi \end{cases}$
(C) $\tan x$
(D) None of these

Question based on

Continuous and discontinuous function

Q.45 Function $f(x) = |x|$ is-

- (A) discontinuous at $x = 0$
(B) discontinuous at $x = 1$
(C) continuous at all point
(D) discontinuous at all points

Q.46 Point of discontinuity for $\sec x$ is -

- (A) $x = -\pi/2$ (B) $x = 3\pi/2$
(C) $x = -5\pi/2$ (D) All of these

Q.47 Function $f(x) = \frac{1}{\log |x|}$ is discontinuous at -

- (A) one point
(B) two points
(C) three points
(D) infinite number of points

Q.48 If $f(x) = x - [x]$, then f is discontinuous at -

- (A) every natural number
(B) every integer
(C) origin
(D) Nowhere

Q.49 Which one is the discontinuous function at any point -

- (A) $\sin x$ (B) x^2
(C) $1/(1-2x)$ (D) $1/(1+x^2)$

Q.50 The point of discontinuity of $\operatorname{cosec} x$ is -

- (A) $x = \pi$ (B) $x = \pi/2$
(C) $x = 3\pi/2$ (D) None of these

Q.51 In the following, continuous function is-

- (A) $\tan x$ (B) $\sec x$
 (C) $\sin 1/x$ (D) None of these
- Q.52** In the following, discontinuous function is-
 (A) $\sin x$ (B) $\cos x$
 (C) $\tan x$ (D) e^x
- Q.53** Which of the following functions is every where continuous-
 (A) $x + |x|$ (B) $x - |x|$
 (C) $x |x|$ (D) All above
- Q.54** Which of the following functions is discontinuous at $x = a$ -
 (A) $\tan(x - a)$ (B) $\sin(x - a)$
 (C) $\operatorname{cosec}(x - a)$ (D) $\sec(x - a)$
- Q.55** If $f(x)$ is continuous and $g(x)$ is discontinuous function, then $f(x) + g(x)$ is-
 (A) continuous function
 (B) discontinuous function
 (C) constant function
 (D) identity function
- Q.56** Function $f(x) = |x-2| - 2|x-4|$ is discontinuous at
 (A) $x = 2, 4$ (B) $x = 2$
 (C) Nowhere (D) Except $x = 2, 4$
- Q.57** Function $f(x) = |\sin x| + |\cos x| + |x|$ is discontinuous at-
 (A) $x = 0$ (B) $x = \pi/2$
 (C) $x = \pi$ (D) No where
- Q.58** Function $f(x) = 1 + |\sin x|$ is-
 (A) continuous only at $x = 0$
 (B) discontinuous at all points
 (C) continuous at all points
 (D) None of these
- Q.59** If function is $f(x) = |x| + |x - 1| + |x - 2|$, then it is -
 (A) discontinuous at $x = 0$
 (B) discontinuous at $x = 0, 1$
 (C) discontinuous at $x = 0, 1, 2$
 (D) everywhere continuous
- Q.60** Function $f(x) = \frac{x^3 - 1}{x^2 - 3x + 2}$ is discontinuous at -
 (A) $x = 1$ (B) $x = 2$
 (C) $x = 1, 2$ (D) No where
- Q.61** If $f(x) = \frac{1}{(1-x)}$ and $g(x) = f[f\{f(x)\}]$, then $g(x)$ is discontinuous at -
 (A) $x = 3$ (B) $x = 2$
 (C) $x = 0$ (D) $x = 4$
- Q.62** The function $f(x) = \frac{|3x-4|}{3x-4}$ is discontinuous at
 (A) $x = 4$ (B) $x = 3/4$
 (C) $x = 4/3$ (D) No where
- Q.63** The function $f(x) = \left(\frac{\pi}{2} - x\right) \tan x$ is discontinuous at-
 (A) $x = \pi$ (B) $x = 0$
 (C) $x = \frac{\pi}{2}$ (D) None of these
- Q.64** Which of the following function has finite number of points of discontinuity-
 (A) $\sin[\pi x]$ (B) $|x|/x$
 (C) $\tan x$ (D) $x + [x]$
- Q.65** The points of discontinuity of
 $f(x) = \tan\left(\frac{\pi x}{x+1}\right)$ other than $x = -1$ are-
 (A) $x = \pi$ (B) $x = 0$
 (C) $x = \frac{2m-1}{2m+1}$
 (D) $x = \frac{2m+1}{1-2m}$, m is any integer
- Q.66** In the following continuous function is-
 (A) $[x]$ (B) $x - [x]$
 (C) $\sin[x]$ (D) $e^x + e^{-x}$
- Q.67** In the following, discontinuous function is-
 (A) $\sin^2 x + \cos^2 x$ (B) $e^x + e^{-x}$
 (C) e^{x^2} (D) $e^{1/x}$

- Q.68** If $f(x)$ is continuous function and $g(x)$ is discontinuous function, then correct statement is -
 (A) $f(x) + g(x)$ is a continuous function
 (B) $f(x) - g(x)$ is a continuous function
 (C) $f(x) + g(x)$ is a discontinuous function
 (D) $f(x)g(x)$ is a continuous function

Question based on

Differentiability of function

- Q.69** Which of the following functions is not differentiable at $x = 0$ -
 (A) $x|x|$ (B) x^3
 (C) e^{-x} (D) $x + |x|$

- Q.70** Which of the following is differentiable function-
 (A) $x^2 \sin \frac{1}{x}$ (B) $x|x|$
 (C) $\cosh x$ (D) all above

- Q.71** The function $f(x) = \sin |x|$ is-
 (A) continuous for all x
 (B) continuous only at certain points
 (C) differentiable at all points
 (D) None of these

- Q.72** If $f(x) = |x-3|$, then f is-
 (A) discontinuous at $x = 2$
 (B) not differentiable at $x = 2$
 (C) differentiable at $x = 3$
 (D) continuous but not differentiable at $x = 3$

- Q.73** If $f(x) = \frac{|x-1|}{x-1}$, $x \neq 1$, and $f(1) = 1$, then the correct statement is-
 (A) discontinuous at $x = 1$
 (B) continuous at $x = 1$
 (C) differentiable at $x = 1$
 (D) discontinuous for $x > 1$

- Q.74** If $f(x) = \begin{cases} x+1, & x > 1 \\ 0, & x = 1 \\ 7-3x, & x < 1 \end{cases}$, then $f'(0)$ equals-
 (A) 1 (B) 2 (C) 0 (D) -3

- Q.75** The function $f(x) = |x| + |x - 1|$ is not differential at -
 (A) $x = 0, 1$ (B) $x = 0, -1$
 (C) $x = -1, 1$ (D) $x = 1, 2$

- Q.76** If $f(x) = \begin{cases} e^{1/x}, & x \neq 0 \\ 0, & x = 0 \end{cases}$, then which one is correct-
 (A) $f(x)$ is differentiable at $x = 0$
 (B) $f(x)$ is discontinuous at $x = 0$
 (C) $f(x)$ is continuous no where
 (D) None of these

- Q.77** Function $[x]$ is not differentiable at -
 (A) every rational number
 (B) every integer
 (C) origin
 (D) every where

- Q.78** If $f(x) = \begin{cases} |x-3|, & \text{when } x \geq 1 \\ \frac{x^2}{4} - \frac{3x}{2} + \frac{13}{4}, & \text{when } x < 1 \end{cases}$, then correct statement is-
 (A) f is discontinuous at $x = 1$
 (B) f is discontinuous at $x = 3$
 (C) f is differentiable at $x = 1$
 (D) f is differentiable at $x = 3$

- Q.79** Function $f(x) = \frac{|x|}{x}$ is-
 (A) continuous every where
 (B) differentiable every where
 (C) differentiable every where except at $x = 0$
 (D) None of these

- Q.80** Let $f(x) = |x-a| + |x-b|$, then-
 (A) $f(x)$ is continuous for all $x \in \mathbb{R}$
 (B) $f(x)$ is differential for $\forall x \in \mathbb{R}$
 (C) $f(x)$ is continuous except at $x = a$ and b
 (D) None of these

- Q.81** Function $f(x) = |x-1| + |x-2|$ is differentiable in $[0, 3]$, except at-
 (A) $x = 0$ and $x = 3$ (B) $x = 1$
 (C) $x = 2$ (D) $x = 1$ and $x = 2$

Q.82 If $f(x) = \begin{cases} 1, & \text{when } x < 0 \\ 1 + \sin x, & \text{when } 0 \leq x \leq \pi/2 \end{cases}$, then at $x = 0$, $f'(x)$ equals-

(A) 1 (B) 0
(C) ∞ (D) Does not exist

Q.83 If $f(x) = \begin{cases} \frac{x}{1+e^{1/x}}, & x \neq 0 \\ 0, & x = 0 \end{cases}$, then the function $f(x)$ is differentiable for -

(A) $x \in \mathbb{R}_+$ (B) $x \in \mathbb{R}$
(C) $x \in \mathbb{R}_0$ (D) None of these

Q.84 If $f(x) = \begin{cases} x^\alpha \sin \frac{1}{x}, & x \neq 0 \\ 0, & x = 0 \end{cases}$ is differentiable at $x = 0$, then-

(A) $\alpha > 0$ (B) $\alpha > 1$
(C) $\alpha \geq 1$ (D) $\alpha \geq 0$

Q.85 If $f(x) = \begin{cases} e^x, & x \leq 0 \\ |1-x|, & x > 0 \end{cases}$, then $f(x)$ is-

(A) continuous at $x = 0$
(B) differentiable at $x = 0$
(C) differentiable at $x = 1$
(D) differentiable both at $x = 0$ and 1

Q.86 The function $f(x) = x - |x|$ is not differentiable at

(A) $x = 1$ (B) $x = -1$
(C) $x = 0$ (D) Nowhere

Q.87 Which of the following function is not differentiable at $x = 1$

(A) $\sin^{-1}x$ (B) $\tan x$
(C) a^x (D) $\sin x$

Q.88 If $f(x) = \begin{cases} \frac{x-1}{2x^2-7x+5}, & x \neq 1 \\ -\frac{1}{3}, & x = 1 \end{cases}$, then $f'(1)$ equals -

(A) $\frac{2}{9}$ (B) $-\frac{2}{9}$
(C) 0 (D) Does not exist

Q.89 If $f(x) = \begin{cases} \frac{\sin x^2}{x}, & x \neq 0 \\ 0, & x = 0 \end{cases}$, then at $x = 0$, $f(x)$ is

(A) continuous and differentiable
(B) neither continuous nor differentiable
(C) continuous but not differentiable
(D) None of these

Q.90 Function $f(x) = 1 + |\sin x|$ is-

(A) continuous no where
(B) differentiable no where
(C) everywhere continuous
(D) None of these

Q.91 Function $f(x) = \begin{cases} x^2, & x \leq 0 \\ 1, & 0 < x \leq 1 \\ 1/x, & x > 1 \end{cases}$ is-

(A) differentiable at $x = 0, 1$
(B) differentiable only at $x = 0$
(C) differentiable at only $x = 1$
(D) Not differentiable at $x = 0, 1$

LEVEL- 2

Q.1 If $[\cdot]$ denotes G.I.F. then, in the following, continuous function is-

- (A) $\cos [x]$ (B) $\sin \pi[x]$
 (C) $\sin \frac{\pi}{2} [x]$ (D) All above

Q.2 If $f(x) = \frac{1 - \cos(1 - \cos x)}{x^4}$, ($x \neq 0$) is continuous

everywhere, then $f(0)$ equals-

- (A) $1/8$ (B) $1/2$
 (C) $1/4$ (D) None of these

Q.3 For function $f(x) = \begin{cases} \left(1 + \frac{4x}{5}\right)^{1/x}, & x \neq 0 \\ e^{4/5}, & x = 0 \end{cases}$, the

correct statement is-

- (A) $f(0+0)$ and $f(0-0)$ do not exist
 (B) $f(0+0) \neq f(0-0)$
 (C) $f(x)$ continuous at $x = 0$
 (D) $\lim_{x \rightarrow 0} f(x) \neq f(0)$

Q.4 If $f(x) = \begin{cases} \frac{\sin(a+1)x + \sin x}{x}, & x < 0 \\ c, & x = 0, \text{ is} \\ \frac{\sqrt{x+bx^2} - \sqrt{x}}{bx\sqrt{x}}, & x > 0 \end{cases}$,

continuous at $x = 0$, then

- (A) $a = 3/2$, $c = 1/2$, b is any real number
 (B) $a = -3/2$, $c = 1/2$, b is $\mathbb{R} - \{0\}$
 (C) $a = 3/2$, $c = -1/2$, $b \in \mathbb{R} - \{0\}$
 (D) None of these

Q.5 Function $f(x) = 4x^3 + 3x^2 + e^{\cos x} + |x-3| + \log(a^x - 1) + x^{1/3}$ ($a > 1$) is discontinuous at-

- (A) $x = 0$ (B) $x = 1$
 (C) $x = 2$ (D) $x = \frac{\pi}{2}$

Q.6 If $f(x) = \frac{\sqrt{a^2 - ax + x^2} - \sqrt{a^2 + ax + x^2}}{\sqrt{a+x} - \sqrt{a-x}}$ is ($a > 0$)

continuous for all values of x , then $f(0)$ is equal to-

- (A) $a\sqrt{a}$ (B) \sqrt{a}
 (C) $-\sqrt{a}$ (D) $-a\sqrt{a}$

Q.7 Function $f(x) = \begin{cases} \frac{(b^2 - a^2)}{2}, & 0 \leq x \leq a \\ \frac{b^2}{2} - \frac{x^2}{6} - \frac{a^3}{3x}, & a < x \leq b, \text{ is} \\ \frac{1}{3} \left(\frac{b^3 - a^3}{x} \right), & x > b \end{cases}$

- (A) continue at $x = a$
 (B) continue at $x = b$
 (C) discontinue on both $x = a, x = b$
 (D) continue at both $x = a, x = b$

Q.8 The function $f(x) = \begin{cases} \frac{e^{1/x} - 1}{e^{1/x} + 1}, & x \neq 0 \\ 0, & x = 0 \end{cases}$,

- (A) is continuous at $x = 0$
 (B) is not continuous at $x = 0$
 (C) is continuous at $x = 2$
 (D) None of these

Q.9 If function $f(x) = \left(\frac{\sin x}{\sin \alpha} \right)^{1/x - \alpha}$ where, $\alpha \neq m\pi$

($m \in \mathbb{I}$) is continuous then -

- (A) $f(\alpha) = e^{\tan \alpha}$ (B) $f(\alpha) = e^{\cot \alpha}$
 (C) $f(\alpha) = e^{2 \cot \alpha}$ (D) $f(\alpha) = \cot \alpha$

Q.10 If $f(x) = \begin{cases} -2 \sin x, & x \leq -\pi/2 \\ a \sin x + b, & -\pi/2 < x < \pi/2, \text{ is a} \\ \cos x, & x \geq \pi/2 \end{cases}$

continuous function for every value x , then-

- (A) $a = b = 1$ (B) $a = b = -1$
 (C) $a = 1, b = -1$ (D) $a = -1, b = 1$

Q.11 If function $f(x) = x - |x - x^2|$, $-1 \leq x \leq 1$ then f is-

- (A) continuous at $x = 0$
- (B) continuous at $x = 1$
- (C) continuous at $x = -1$
- (D) everywhere continuous

Q.12 $f(x) = 1 + 2^{1/x}$ is-

- (A) continuous everywhere
- (B) continuous nowhere
- (C) discontinuous at $x = 0$
- (D) None of these

Q.13 Let $[.]$ denotes G.I.F. and $f(x) = [x] + [-x]$ and m is any integer, then correct statement is -

- (A) $\lim_{x \rightarrow m} f(x)$ does not exist
- (B) $f(x)$ is continuous at $x = m$
- (C) $\lim_{x \rightarrow m} f(x)$ exists
- (D) None of these

Q.14 If $f(x) = (\tan x \cot \alpha)^{1/(x-\alpha)}$ is continuous at $x = \alpha$, then the value of $f(\alpha)$ is -

- (A) $e^{2 \sin 2\alpha}$
- (B) $e^{2 \operatorname{cosec} 2 \alpha}$
- (C) $e^{\operatorname{cosec} 2 \alpha}$
- (D) $e^{\sin 2 \alpha}$

Q.15 Let $[.]$ denotes G.I.F. for the function

$$f(x) = \frac{\tan(\pi[x - \pi])}{1 + [x]^2}$$

the wrong statement is -

- (A) $f(x)$ is discontinuous at $x = 0$
- (B) $f(x)$ is continuous for all values of x
- (C) $f(x)$ is continuous at $x = 0$
- (D) $f(x)$ is a constant function

Q.16 The point of discontinuity of the function

$$f(x) = \frac{1 + \cos 5x}{1 - \cos 4x}$$

is-

- (A) $x = 0$
- (B) $x = \pi$
- (C) $x = \pi/2$
- (D) All the above

Q.17 Let $f(x) = \frac{\sqrt{1 + \sin x} - \sqrt{1 - \sin x}}{x}$. The value

which should be assigned to f at $x = 0$ so that it is continuous everywhere is-

- (A) 1
- (B) 2
- (C) -2
- (D) 1/2

Q.18 If the function

$$f(x) = \begin{cases} \frac{\sin(k+1)x + \sin x}{x}, & \text{when } x < 0 \\ 1/2, & \text{when } x = 0 \\ \frac{(x + 2x^2)^{1/2}}{2x^{3/2}}, & \text{when } x > 0 \end{cases}$$

continuous at $x = 0$, then the value of k is-

- (A) 1/2
- (B) -1/2
- (C) -3/2
- (D) 1

Q.19 If $f(x) = \begin{cases} 3, & x < 0 \\ 2x + 1, & x \geq 0 \end{cases}$ then -

- (A) both $f(x)$ and $f(|x|)$ are differentiable at $x = 0$
- (B) $f(|x|)$ is differentiable but $f(x)$ is not differentiable at $x = 0$
- (C) $f(x)$ is differentiable but $f(|x|)$ is not differentiable at $x = 0$
- (D) both $f(x)$ and $f(|x|)$ are not differentiable at $x = 0$

Q.20 The number of points in the interval $(0, 2)$ where the derivative of the function

$f(x) = |x - 1/2| + |x - 1| + \tan x$ does not exist is-

- (A) 1
- (B) 2
- (C) 3
- (D) 4

Q.21 Function $f(x) = \sin(\pi[x])$ is-

- (A) differentiable every where
- (B) differentiable no where
- (C) not differentiable at $x = 1$ and -1
- (D) None of these

Q.22 Function $f(x) = \begin{cases} x \tan^{-1}(1/x), & x \neq 0 \\ 0, & x = 0 \end{cases}$ at $x = 0$

is-

- (A) discontinuous
- (B) continuous
- (C) differentiable
- (D) None of these

Q.23 Function $f(x) = \frac{\cos x - \sin x}{\sin 4x}$ is not defined at

$x = \frac{\pi}{4}$. The value which should be assigned to

f at $x = \frac{\pi}{4}$, so that it is continuous there, is-

- (A) 0
- (B) $\frac{1}{2\sqrt{2}}$
- (C) $-\frac{1}{\sqrt{2}}$
- (D) None

- Q.24** Let $f(x) = \max \{2 \sin x, 1 - \cos x\}$, $x \in (0, \pi)$.
Then set of points of non-differentiability is -
(A) ϕ (B) $\{\pi/2\}$
(C) $\{\pi - \cos^{-1} 3/5\}$ (D) $\{\cos^{-1} 3/5\}$

Q.25 If $f(x) = \begin{cases} x \frac{e^{1/x} - e^{-1/x}}{e^{1/x} + e^{-1/x}}, & x \neq 0 \\ 0, & x = 0 \end{cases}$, then correct

statement is-

- (A) f is continuous at all points except $x = 0$
(B) f is continuous at every point but not differentiable
(C) f is differentiable at every point
(D) f is differentiable only at the origin

Q.26 Consider the following statements-

- (I) If a function is differentiable at some point then it must be continuous at that point
(II) If a function continuous at some point then it is not necessary that it is differentiable at that point.
(III) Differentiability of a function at some point is the necessary and sufficient condition for continuity at that point.

From above, correct statements are-

- (A) I, II, III (B) I, III
(C) I, II (D) II, III

Q.27 State which of the following is a false statement -

- (A) If $f(x)$ is continuous at $x = a$ then

$$f(a) = \lim_{x \rightarrow a} f(x)$$

- (B) If $\lim_{x \rightarrow a} f(x)$ exists, then $f(x)$ is continuous at

$$x = a$$

- (C) If $f(x)$ is differentiable at $x = a$, then it is continuous at $x = a$

- (D) If $f(x)$ is continuous at $x = a$, then $\lim_{x \rightarrow a} f(x)$ exists

LEVEL- 3

Q.1 If the derivative of the function -

$$f(x) = \begin{cases} ax^2 + b, & x < -1 \\ bx^2 + ax + 4, & x \geq -1 \end{cases}$$

is everywhere continuous, then

- (A) $a = 2, b = 3$ (B) $a = 3, b = 2$
 (C) $a = -2, b = -3$ (D) $a = -3, b = -2$

Q.2 The value of $f(0)$, so that the function

$$f(x) = \frac{(27 - 2x)^{1/3} - 3}{9 - 3(243 + 5x)^{1/5}}, \quad (x \neq 0)$$
 is continuous,

is given by -

- (A) $2/3$ (B) 6 (C) 2 (D) 4

Q.3 If $f(x) = \begin{cases} |x-4|, & \text{for } x \geq 1 \\ (x^3/2) - x^2 + 3x + (1/2), & \text{for } x < 1 \end{cases}$, then

- (A) $f(x)$ is continuous at $x = 1$ and at $x = 4$
 (B) $f(x)$ is differentiable at $x = 4$
 (C) $f(x)$ is continuous and differentiable at $x = 1$
 (D) $f(x)$ is only continuous at $x = 1$

Q.4 Let $f(x) = |x|$ and $g(x) = |x^3|$, then -

- (A) $f(x)$ & $g(x)$ both are continuous at $x = 0$
 (B) $f(x)$ & $g(x)$ both are differentiable at $x = 0$
 (C) $f(x)$ is differentiable but $g(x)$ is not differentiable at $x = 0$
 (D) $f(x)$ & $g(x)$ both are not differentiable at $x = 0$

Q.5 Let $f(x) = \begin{cases} x^n \sin \frac{1}{x}, & x \neq 0 \\ 0, & x = 0 \end{cases}$. Then $f(x)$ is

continuous but not differentiable at $x = 0$ if -

- (A) $n \in (0, 1]$ (B) $n \in [0, \infty)$
 (C) $n \in (-\infty, 0)$ (D) $n = 0$

Q.6 If $f(x) = a |\sin x| + b e^{|x|} + c |x|^3$ and if $f(x)$ is differentiable at $x = 0$, then -

- (A) $a = b = c = 0$
 (B) $a = 0, b = 0; c \in \mathbb{R}$
 (C) $b = c = 0; a \in \mathbb{R}$
 (D) $c = 0, a = 0; b \in \mathbb{R}$

Q.7 The set of points where function

$$f(x) = \sqrt{1 - e^{-x^2}}$$
 is differentiable is -

- (A) $(-\infty, \infty)$ (B) $(-\infty, 0) \cup (0, \infty)$
 (C) $(-1, \infty)$ (D) none of these

Q.8 Let $f(x) = \begin{cases} \sin 2x, & 0 < x \leq \pi/6 \\ ax + b, & \pi/6 < x < 1 \end{cases}$; If $f(x)$ and

$f'(x)$ are continuous, then -

- (A) $a = 1, b = \frac{1}{\sqrt{2}} + \frac{\pi}{6}$
 (B) $a = \frac{1}{\sqrt{2}}, b = \frac{1}{\sqrt{2}}$
 (C) $a = 1, b = \frac{\sqrt{3}}{2} - \frac{\pi}{6}$
 (D) none of these

Q.9 Let $f(x) = \lim_{n \rightarrow \infty} (\sin x)^{2n}$; then f is -

- (A) discontinuous at $x = 3\pi/2$
 (B) discontinuous at $x = \pi/2$
 (C) discontinuous at $x = -\pi/2$
 (D) All the above

Q.10 Let $[.]$ denotes G.I.F. and if function

$$f(x) = \left(\frac{x}{2} - 1 \right)$$
 then in the interval $[0, \pi]$

- (A) $\tan [f(x)]$ is discontinuous but $1/f(x)$ is continuous
 (B) $\tan [f(x)]$ is continuous but $\frac{1}{f(x)}$ is discontinuous
 (C) $\tan [f(x)]$ and $f^{-1}(x)$ is continuous
 (D) $\tan [f(x)]$ and $1/f(x)$ both are discontinuous

Q.11 The function $f(x) = \frac{4 - x^2}{4x - x^3}$ is equal to -

- (A) discontinuous at only one point
 (B) discontinuous exactly at two points
 (C) discontinuous exactly at three points
 (D) none of these

- Q.12** The function $f(x) = \sin^{-1}(\cos x)$ is -
 (A) discontinuous at $x = 0$
 (B) continuous at $x = 0$
 (C) differentiable at $x = 0$
 (D) none of these
- Q.13** The function $f(x) = e^{-|x|}$ is -
 (A) continuous everywhere but not differentiable at $x = 0$
 (B) continuous and differentiable everywhere
 (C) not continuous at $x = 0$
 (D) none of these
- Q.14** If $x + 4|y| = 6y$, then y as a function of x is -
 (A) continuous at $x = 0$ (B) derivable at $x = 0$
 (C) $\frac{dy}{dx} = \frac{1}{2}$ for all x (D) none of these
- Q.15** Let $f(x + y) = f(x) + f(y)$ and $f(x) = x^2 g(x)$ for all $x, y, \in \mathbb{R}$, where $g(x)$ is continuous function. Then $f'(x)$ is equal to -
 (A) g' (B) $g(x)$
 (C) $f(x)$ (D) none of these
- Q.16** Let $f(x + y) = f(x) f(y)$ for all $x, y, \in \mathbb{R}$, Suppose that $f(3) = 3$ and $f'(0) = 11$ then $f'(3)$ is equal to-
 (A) 22 (B) 44
 (C) 28 (D) none of these

➤ Statement type Questions

All questions are Assertion & Reason type questions. Each of these questions contains two statements: Statement-1 (Assertion) and Statement-2 (Reason). Answer these questions from the following four option.

- (A) Statement-I and Statement-II are true Statement-II is the correct explanation of Statement-I
 (B) Statement-I Statement-II are true but Statement-II is not the correct explanation of Statement-I.
 (C) Statement-I is true but Statement-II is false
 (D) Statement-I is false but Statement-II is true.

Q.17 **Statement-1 :**

$f(x) = \frac{1}{x - [x]}$ is discontinuous for integral values of x

Statement-2 : For integral values of x , $f(x)$ is undefined.

Q.18 **Statement-1 :**

If $f(x) = \frac{(e^{kx} - 1)\sin kx}{4x^2}$ ($x \neq 0$) and $f(0) = 9$ is

continuous at $x = 0$ then $k = \pm 6$.

Statement-2 : For continuous function

$$\lim_{x \rightarrow 0} f(x) = f(0)$$

Q.19 **Statement I:**

$y = \frac{x}{1 + |x|}$, $x \in \mathbb{R}$, $f(x)$ is differentiable

every where.

Statement II :

$$f(x) = \frac{x}{1 + |x|}, x \in \mathbb{R} \text{ then } f'(x) = \begin{cases} \frac{1}{(1+x)^2}, x \geq 0 \\ \frac{1}{(1-x)^2}, x < 0 \end{cases}$$

Q.20 **Statement-1 :** If $f(x) = \lim_{n \rightarrow \infty} (\sin x)^{2n}$, then the

set of points discontinuities of f is

$$\left\{ (2n+1)\frac{\pi}{2}, n \in \mathbb{I} \right\}$$

Statement-2 : Since $-1 < \sin x < 1$, as $n \rightarrow \infty$, $(\sin x)^{2n} \rightarrow 0$, $\sin x = \pm 1 \Rightarrow \pm (1)^{2n} \rightarrow 1, n \rightarrow \infty$

Q.21 **Statement I :**

$f(x) = |x - 2|$ is differentiable at $x = 2$.

Statement II :

$f(x) = |x - 2|$ is continuous at $x = 2$.

Q.22 **Statement-1 :** The function

$y = \sin^{-1}(\cos x)$ is not differentiable at

$x = n\pi, n \in \mathbb{Z}$ is particular at $x = \pi$

Statement-2 : $\frac{dy}{dx} = \frac{-\sin x}{|\sin x|}$ so the function is

not differentiable at the points where $\sin x = 0$.

Q.23 **Statement-1 :**

The function $f(x) = |x^3|$ is differentiable at $x = 0$

Statement-2 : at $x = 0$, $f'(x) = 0$

Q.24 **Statement I :** $f(x) = \sin x$ and $g(x) = \operatorname{sgn}(x)$ then $f(x)g(x)$ is differentiable at $x = 1$.

Statement II : Product of two differentiable function is differentiable function

➤ **Passage Based Questions**

$$\text{Let } f(x) = \begin{cases} \frac{a(1 - x \sin x) + b \cos x + 5}{x^2}, & x < 0 \\ \frac{3}{3}, & x = 0 \\ \left\{ 1 + \left(\frac{cx + dx^3}{x^2} \right) \right\}^{1/x}, & x > 0 \end{cases}$$

If f is continuous at $x = 0$

On the basis of above information, answer the following questions :-

- Q.25** The value of a is -
 (A) -1 (B) $\ln 3$ (C) 0 (D) -4
- Q.26** The value of b is -
 (A) -1 (B) $\ln 3$ (C) 0 (D) -4
- Q.27** The value of c is
 (A) 2 (B) 3
 (C) 0 (D) none of these
- Q.28** The value of e^d is -
 (A) 0 (B) 1 (C) 2 (D) 3

➤ **Column Matching Questions**

Match the entry in Column I with the entry in Column II.

- | Q.29 | Column-I | Column-II |
|-------------|---|---|
| (A) | $f(x) = x^2 \sin(1/x), x \neq 0$
$f(0) = 0$ | (P) continuous but not derivable |
| (B) | $f(x) = \frac{1}{1 - e^{-1/x}}, x \neq 0$,
and $f(0) = 0$ | (Q) f is differentiable
f' is not continuous |
| (C) | $f(x) = x \sin 1/x, x \neq 0$
$f(0) = 0$ | (R) f is not continuous |
| (D) | $f(x) = x^3 \sin 1/x, x \neq 0$
$f(0) = 0$ | (S) f' is continuous but not derivable |
-
- | Q.30 | Column I | Column II |
|-------------|--------------------------|--|
| (A) | $f(x) = x^3 $ is | (P) continuous in $(-1, 1)$ |
| (B) | $f(x) = \sqrt{ x }$ | (Q) differentiable in $(-1, 1)$ |
| (C) | $f(x) = \sin^{-1}x $ is | (R) differentiable in $(0, 1)$ |
| (D) | $f(x) = x $ is | (S) not differentiable
at least at one point in $(-1, 1)$ |

LEVEL- 4

(Question asked in previous AIEEE and IIT-JEE)

SECTION –A

- Q.1** If $f(x) = \begin{cases} x & x \in \mathbb{Q} \\ -x & x \notin \mathbb{Q} \end{cases}$, then f is continuous at-
- [AIEEE-2002]
- (A) only at zero
(B) only at 0, 1
(C) all real numbers
(D) all rational numbers
- Q.2** If for all values of x & y ; $f(x + y) = f(x) \cdot f(y)$ and $f(5) = 2$ $f'(0) = 3$, then $f'(5)$ is-
- [AIEEE- 2002]
- (A) 3 (B) 4
(C) 5 (D) 6
- Q.3** If $f(x) = \begin{cases} x e^{-\left(\frac{1}{|x|} + \frac{1}{x}\right)}, & x \neq 0 \\ 0, & x = 0 \end{cases}$ then $f(x)$ is
- [AIEEE- 2003]
- (A) discontinuous everywhere
(B) continuous as well as differentiable for all x
(C) continuous for all x but not differentiable at $x = 0$
(D) neither differentiable nor continuous at $x = 0$
- Q.4** Let $f(x) = \frac{1 - \tan x}{4x - \pi}$, $x \neq \frac{\pi}{4}$, $x \in \left[0, \frac{\pi}{2}\right]$. If $f(x)$ is continuous in $\left[0, \frac{\pi}{2}\right]$, then $f\left(\frac{\pi}{4}\right)$ is-
- [AIEEE- 2004]
- (A) 1 (B) 1/2
(C) -1/2 (D) -1
- Q.5** If f is a real-valued differentiable function satisfying $|f(x) - f(y)| \leq (x - y)^2$, $x, y \in \mathbb{R}$ and $f(0) = 0$, then $f(1)$ equals-
- [AIEEE-2005]
- (A) -1 (B) 0
(C) 2 (D) 1
- Q.6** Suppose $f(x)$ is differentiable at $x = 1$ and $\lim_{h \rightarrow 0} \frac{1}{h} f(1 + h) = 5$, then $f'(1)$ equals -
- [AIEEE-2005]
- (A) 3 (B) 4 (C) 5 (D) 6
- Q.7** The set of points where $f(x) = \frac{x}{1 + |x|}$ is differentiable is -
- [AIEEE- 2006]
- (A) $(-\infty, -1) \cup (-1, \infty)$
(B) $(-\infty, \infty)$
(C) $(0, \infty)$
(D) $(-\infty, 0) \cup (0, \infty)$
- Q.8** The function $f : \mathbb{R} \setminus \{0\} \rightarrow \mathbb{R}$ given by $f(x) = \frac{1}{x} - \frac{2}{e^{2x} - 1}$ can be made continuous at $x = 0$ by defining $f(0)$ as -
- [AIEEE- 2007]
- (A) 2 (B) -1
(C) 0 (D) 1
- Q.9** Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a function defined by $f(x) = \text{Min} \{x + 1, |x| + 1\}$. Then which of the following is true?
- [AIEEE 2007]
- (A) $f(x) \geq 1$ for all $x \in \mathbb{R}$
(B) $f(x)$ is not differentiable at $x = 1$
(C) $f(x)$ is differentiable everywhere
(D) $f(x)$ is not differentiable at $x = 0$
- Q.10** Let $f(x) = \begin{cases} (x-1) \sin \frac{1}{x-1}, & \text{if } x \neq 1 \\ 0, & \text{if } x = 1 \end{cases}$
- Then which one of the following is true?
- [AIEEE 2008]
- (A) f is differentiable at $x = 0$ and at $x = 1$
(B) f is differentiable at $x = 0$ but not at $x = 1$
(C) f is differentiable at $x = 1$ but not at $x = 0$
(D) f is neither differentiable at $x = 0$ nor at $x = 1$

- Q.4** The function $f(x) = [x] \cos \{(2x - 1)/2\}\pi$, $[\]$ denotes the greatest integer function, is discontinuous at **[IIT-1995]**
 (A) all x
 (B) all integer points
 (C) no x
 (D) x which is not an integer
- Q.5** Let $f(x)$ be defined for all $x > 0$ and be continuous. Let $f(x)$ satisfy $f\left(\frac{x}{y}\right) = f(x) - f(y)$ for all x, y & $f(e) = 1$. Then- **[IIT Scr.95]**
 (A) $f(x)$ is bounded
 (B) $f\left(\frac{1}{x}\right) \rightarrow 0$ as $x \rightarrow 0$
 (C) $x f(x) \rightarrow 1$ as $x \rightarrow 0$
 (D) $f(x) = \log x$
- Q.6** The function $f(x) = [x]^2 - [x^2]$ (where $[y]$ is the greatest integer less than or equal to y), is discontinuous at - **[IIT-1999]**
 (A) All integers
 (B) All integers except 0 and 1
 (C) All integers except 0
 (D) All integers except 1
- Q.7** Indicate the correct alternative:
 Let $[x]$ denote the greater integer $\leq x$ and $f(x) = [\tan^2 x]$, then **[IIT-1993]**
 (A) $\lim_{x \rightarrow 0} f(x)$ does not exist
 (B) $f(x)$ is continuous at $x = 0$
 (C) $f(x)$ is not differentiable at $x = 0$
 (D) $f'(0) = 1$
- Q.8** $g(x) = x f(x)$, where $f(x) = \begin{cases} x \sin(1/x), & x \neq 0 \\ 0 & x = 0 \end{cases}$ at $x = 0$ **[IIT-1994]**
 (A) g is differentiable but g' is not continuous
 (B) both f and g are differentiable
 (C) g is differentiable but g' is continuous
 (D) None of these
- Q.9** Let $f\left(\frac{x+y}{2}\right) = \frac{f(x)+f(y)}{2}$ for all real x and y and $f'(0) = -1$, $f(0) = 1$, then $f'(2) =$ **[IIT-1995]**
 (A) $1/2$ (B) 1 (C) -1 (D) $-1/2$
- Q.10** Let $h(x) = \min\{x, x^2\}$, for every real number of x . Then - **[IIT-1998]**
 (A) h is not differentiable at two values of x
 (B) h is differentiable for all x
 (C) $h'(x) = 0$, for all $x > 1$
 (D) None of these
- Q.11** The function $f(x) = (x^2 - 1)|x^2 - 3x + 2| + \cos(|x|)$ is not differentiable at. **[IIT-1999]**
 (A) -1 (B) 0 (C) 1 (D) 2
- Q.12** Let $f: \mathbb{R} \rightarrow \mathbb{R}$ is a function which is defined by $f(x) = \max\{x, x^3\}$ set of points on which $f(x)$ is not differentiable is **[IIT Scr. 2001]**
 (A) $\{-1, 1\}$ (B) $\{-1, 0\}$
 (C) $\{0, 1\}$ (D) $\{-1, 0, 1\}$
- Q.13** Find left hand derivative at $x = k$, $k \in \mathbb{I}$.
 $f(x) = [x] \sin(\pi x)$ **[IIT Scr. 2001]**
 (A) $(-1)^k (k-1)\pi$ (B) $(-1)^{k-1} (k-1)\pi$
 (C) $(-1)^k (k-1)k\pi$ (D) $(-1)^{k-1} (k-1)k\pi$
- Q.14** Which of the following functions is differentiable at $x = 0$? **[IIT Scr. 2001]**
 (A) $\cos(|x|) + |x|$ (B) $\cos(|x|) - |x|$
 (C) $\sin(|x|) + |x|$ (D) $\sin(|x|) - |x|$
- Q.15** $f(x) = ||x| - 1|$ is not differentiable at $x =$ **[IIT Scr.2005]**
 (A) $0, \pm 1$ (B) ± 1
 (C) 0 (D) 1
- Q.16** Let $g(x) = \frac{(x-1)^n}{\log \cos^m(x-1)}$; $0 < x < 2$, m and n are integers, $m \neq 0$, $n > 0$, and let p be the left hand derivative of $|x - 1|$ at $x = 1$. If $\lim_{x \rightarrow 1^+} g(x) = p$, then **[IIT- 2008]**
 (A) $n = 1, m = 1$ (B) $n = 1, m = -1$
 (C) $n = 2, m = 2$ (D) $n > 2, m = n$

Q.17 Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a function such that
 $f(x+y) = f(x) + f(y), \forall x, y \in \mathbb{R}$
 If $f(x)$ is differentiable at $x = 0$, then

[IIT- 2011]

- (A) $f(x)$ is differentiable only in a finite interval containing zero
- (B) $f(x)$ is continuous $\forall x \in \mathbb{R}$
- (C) $f'(x)$ is constant $\forall x \in \mathbb{R}$
- (D) $f(x)$ is differentiable except at finitely many points

Q.18 If

$$f(x) = \begin{cases} -x - \frac{\pi}{2}, & x \leq -\frac{\pi}{2} \\ -\cos x, & -\frac{\pi}{2} < x \leq 0 \\ x - 1, & 0 < x \leq 1 \\ \ln x, & x > 1 \end{cases}, \text{ then}$$

[IIT- 2011]

- (A) $f(x)$ is continuous at $x = -\frac{\pi}{2}$
- (B) $f(x)$ is not differentiable at $x = 0$
- (C) $f(x)$ is differentiable at $x = 1$
- (D) $f(x)$ is differentiable at $x = -\frac{3}{2}$

Q.19 For every integer n , let a_n and b_n be real numbers.

Let function $f: \mathbb{R} \rightarrow \mathbb{R}$ be given by

$$f(x) = \begin{cases} a_n + \sin \pi x, & \text{for } x \in [2n, 2n+1] \\ b_n + \cos \pi x, & \text{for } x \in (2n-1, 2n) \end{cases}, \text{ for}$$

all integers n . If f is continuous, then which of the following hold(s) for all n ? [IIT- 2012]

- (A) $a_{n-1} - b_{n-1} = 0$
- (B) $a_n - b_n = 1$
- (C) $a_n - b_{n+1} = 1$
- (D) $a_{n-1} - b_n = -1$

Q.20 Let $f(x) = \begin{cases} x^2 \left| \cos \frac{\pi}{x} \right|, & x \neq 0 \\ 0, & x = 0 \end{cases}, x \in \mathbb{R},$

then f is

[IIT-2012]

- (A) differentiable both at $x = 0$ and at $x = 2$
- (B) differentiable at $x = 0$ but not differentiable at $x = 2$
- (C) not differentiable at $x = 0$ but differentiable at $x = 2$
- (D) differentiable neither at $x = 0$ nor at $x = 2$

ANSWER KEY

LEVEL-1

Que	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
Ans.	D	C	A	D	C	A	C	B	D	D	D	D	A	C	D	A	D	C	A	C
Que	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40
Ans.	B	C	D	B	D	B	D	A	A	A	D	B	A	C	C	A	C	D	D	C
Que	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60
Ans.	C	D	C	C	C	D	C	B	C	A	D	C	D	C	B	C	D	C	D	C
Que	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80
Ans.	C	C	C	B	D	D	D	C	D	D	A	D	A	D	A	B	B	C	C	A
Que	81	82	83	84	85	86	87	88	89	90	91									
Ans.	D	D	C	B	A	C	A	B	A	C	D									

LEVEL-2

Que	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
Ans.	B	A	C	B	A	C	D	B	B	D	D	C	C	B	A	D	A	C	D	C
Que	21	22	23	24	25	26	27													
Ans	A	B	B	C	B	C	B													

LEVEL-3

Que	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
Ans.	A	C	A	A	A	B	B	C	D	D	C	B	A	A	D	D	A	A	A	A
Que	21	22	23	24	25	26	27	28												
Ans	D	A	A	A	A	D	C	D												

29. $A \rightarrow Q$; $B \rightarrow R$; $C \rightarrow P$; $D \rightarrow S$

30. $A \rightarrow P, Q, R$; $B \rightarrow P, R, S$; $C \rightarrow P, R, S$; $D \rightarrow P, R, S$

LEVEL- 4

SECTION-A

Que	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Ans.	A	D	C	C	B	C	B	D	C	B	C	A	C	D	B

SECTION-B

$$1.[A] \quad f(x) = \begin{cases} \frac{1 - \cos 4x}{x^2} & : x < 0 \\ a & : x = 0 \\ \frac{\sqrt{x}}{\sqrt{16 + \sqrt{x}} - 4} & : x > 0 \end{cases}$$

$\therefore f(x)$ is continuous at $x = 0$

$$\therefore \text{R.H.L.} \lim_{h \rightarrow 0} \frac{1 - \cos(4(0-h))}{(0-h)^2}$$

$$\lim_{h \rightarrow 0} \frac{1 - \cos 4h}{h^2} \quad \left(\frac{0}{0} \text{ form} \right)$$

$$= \lim_{h \rightarrow 0} \frac{0 + 4 \sin 4h}{2h} \times \frac{2}{2}$$

$$= 8$$

$$\therefore \text{L.H.L.} = f(0)$$

$$\Rightarrow 8 = a$$

2.[C]

(A) $\tan x$ is discontinuous at $\pi/2$ in $(0, \pi)$

$$(B) \quad f(x) = \begin{cases} x \sin x & ; 0 < x \leq \pi/2 \\ \frac{\pi}{2} \sin(\pi + x) & ; \frac{\pi}{2} < x < \pi \end{cases}$$

$$\text{at } x = \frac{\pi}{2}$$

$$\text{L.H.L.} \lim_{h \rightarrow 0} \left(\frac{\pi}{2} - h \right) \sin \left(\frac{\pi}{2} - h \right)$$

$$= \pi/2 \sin \pi/2 = \pi/2$$

$$\text{R.H.L.} \lim_{h \rightarrow 0} \pi/2 \sin (\pi + \pi/2 + h)$$

$$= \pi/2 \sin 3\pi/2$$

$$= -\pi/2$$

L.H.L. \neq R.H.L.

$\therefore f(x)$ is not continuous at $x = \pi/4$

$$(C) \quad f(x) = \begin{cases} 1, & 0 < x \leq \frac{3\pi}{4} \\ 2\sin \frac{2}{9}x, & \frac{3\pi}{4} < x < \pi \end{cases}$$

at $x = 3\pi/4$ L.H.L. = 1

$f(3\pi/4) = 1$

R.H.L. $\lim_{h \rightarrow 0} 2\sin \frac{2}{9}(3\pi/4 + h)$

$2\sin \frac{2}{9} \frac{3\pi}{4}$

$= 2\sin \pi/6 = 2 \times \frac{1}{2} = 1$

$\therefore f(x)$ is continuous at $x = \frac{3\pi}{4}$

$$3.[A] \quad f(x) = \begin{cases} x \sin x; & 0 < x \leq \pi/2 \\ \frac{\pi}{2} \sin(\pi + x); & \frac{\pi}{2} < x < \pi \end{cases}$$

at $x = \frac{\pi}{2}$

L.H.L. $\lim_{h \rightarrow 0} \left(\frac{\pi}{2} - h\right) \sin\left(\frac{\pi}{2} - h\right)$

$= \pi/2 \sin \pi/2 = \pi/2$

R.H.L. $\lim_{h \rightarrow 0} \pi/2 \sin(\pi + \pi/2 + h)$

$= \pi/2 \sin 3\pi/2$

$= -\pi/2$

L.H.L. \neq R.H.L.

$\therefore f(x)$ is not continuous at $x = \pi/4$

$$4.[C] \quad f(x) = [x] \cos(2x-1) \times \pi/2$$

let $x = n, n \in I$

$f(n) = n \cos(2n-1) \pi/2 = 0$

$f(n+) = n \cos(2n-1) \pi/2 = 0$

$f(n-) = (n-1) \cos(2n-1) \pi/2 = 0$

$$\left(\because \cos(2n-1) \frac{\pi}{2} = 0\right)$$

continuous for all x .

$$5.[D] \quad f(x) = k \ell n x$$

put $x = e$

$k = 1$

$\therefore f(x) = \ell n x$

$$6.[D] \quad f(x) = [x]^2 - [x^2]$$

Let us check continuity at $x = 0$ & 1

L.H.L. $\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} [x]^2 - [x^2]$

$= \lim_{h \rightarrow 0} [0-h]^2 - [(0-h)^2]$

$= +1 - 0 = 1$

R.H.L. $\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} [x]^2 - [x^2]$

$= \lim_{h \rightarrow 0} [0+h]^2 - [(0+h)^2]$

$0 - 0 = 0$

L.H.L. \neq R.H.L.

$\therefore f(x)$ is not continuous at $x = 0$

at $x = 1$

L.H.L. $\lim_{x \rightarrow 1^-} f(x) = \lim_{h \rightarrow 0^-} [1-h]^2 - [(1-h)^2]$

$0 - 0 = 0$

R.H.L. $\lim_{x \rightarrow 1^+} f(x) = \lim_{h \rightarrow 0} [(1+h)^2] - [(1+h)^2]$

$= 1 - 1 = 0$

$f(1) = [1]^2 - [1^2] = 0$

$\therefore f(x)$ is continuous at $x = 1$

clearly $f(x)$ is discontinuous at all other integers except 1

$$7.[B] \quad f(x) = [\tan^2 x]$$

$\lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} [\tan^2 x]$

for continuity at $x = 0$

$f(0) = [\tan^2 0] = [0] = 0$

L.H.L. $\lim_{h \rightarrow 0} [\tan^2(0-h)]$

$= \lim_{h \rightarrow 0} [\tan^2 h] = [\text{Value greater than 0 less than 1}]$

$= 0$

R.H.L. $\lim_{h \rightarrow 0} [\tan^2(0+h)]$

$= \lim_{h \rightarrow 0} [\tan^2 h]$

$= [\text{Value greater than 0 & less than 1}] = 0$

\therefore L.H.L. = R.H.L. = $f(0)$

$\therefore f(x)$ is continuous at $x = 0$

$$8.[A] \quad g(x) = x f(x) \text{ \&}$$

$$f(x) = \begin{cases} x \sin 1/x, & x \neq 0 \\ 0 & x = 0 \end{cases}$$

$$g(x) = x f(x) = \begin{cases} x^2 \sin \frac{1}{x} : x \neq 0 \\ 0 & x = 0 \end{cases}$$

\therefore we know function $\begin{cases} x^\alpha \sin \frac{1}{x} : x \neq 0 \\ 0 & x = 0 \end{cases}$ is

differentiable when $\alpha > 1$

in $g(x) \alpha = 2 \therefore$ it is differentiable

Now $g'(x) = \begin{cases} x^2 \cos \frac{1}{x} \left(-\frac{1}{x^2}\right) + \sin \frac{1}{x} \cdot 2x; & x \neq 0 \\ 0 & x = 0 \end{cases}$

$= \begin{cases} -\cos \frac{1}{x} + 2x \sin \frac{1}{x}; & x \neq 0 \\ 0 & x = 0 \end{cases}$

for continuity at $x = 0$


L.H.L. $\lim_{h \rightarrow 0} -\cos \frac{1}{0-h} + 2(0-h) \sin \frac{1}{0-h}$
 $\lim_{h \rightarrow 0} -\cos \frac{1}{h} + 2h \sin \frac{1}{h}$
 $-\cos \frac{1}{0} + 2 \times 0 = -$ (value between -1 and 1) \neq unique
 $\Rightarrow g'(x)$ is discontinuous at $x = 0$

9.[C] $f\left(\frac{x+y}{2}\right) = \frac{f(x)+f(y)}{2}$
 differentiating both side keeping y as constant

$f\left(\frac{x+y}{2}\right) \left[\frac{1+0}{2}\right] = \frac{f'(x)+0}{2}$
 $\Rightarrow \frac{1}{2} f' \left(\frac{x+y}{2}\right) = \frac{f'(x)}{2}$

put $x = 0$
 $f'(y/2) = -1$
 put $y = 4$
 $f'(2) = -1$

10.[A]



minimum (x, x^2)
 sharp point at $x = 0, 1$
 \Rightarrow Not differentiable at $x = 0, 1$


11.[D]

$(x+1)(x-1) | (x-1)(x-2) | + \cos x$
 $\downarrow \quad \downarrow \quad \downarrow \quad \downarrow$
 D D N.D at $x = 2$ D

$(\because \cos |x| = \cos x)$

\Rightarrow Not differentiable at $x = 2$

12.[D] $f(x) = \max. \{x, x^3\}$
 by graph



$$\therefore f(x) = \begin{cases} x & ; x \leq -1 \\ x^3 & ; -1 \leq x \leq 0 \\ x & ; 0 \leq x \leq 1 \\ x^3 & ; x \geq 1 \end{cases}$$

at $x = 1, -1, 0$ there is sharp point
 $\therefore f(x)$ is not differentiable at these points

13.[A] $f(x) = [x] \sin \pi x$
 at $x = k$ we have to find out L.H.D.
 $\therefore x$ is just less than k
 $\therefore [k-h] = k-1$
 $\therefore f(x) = (k-1) \sin \pi x$

L.H.L. of $f(x) = \lim_{x \rightarrow k^-} \frac{f(x)-f(k)}{x-k}$
 $= \lim_{h \rightarrow 0} \frac{f(k-h)-f(k)}{-h}$
 $= \lim_{h \rightarrow 0} \frac{(k-1) \sin(\pi)(k-h) - k \sin \pi k}{-h} : |k \in \mathbb{I}|$
 $= \lim_{h \rightarrow 0} \frac{(k-1) \sin(\pi k - \pi h) - 0}{-h}$
 $= \lim_{h \rightarrow 0} \frac{(k-1)(-1)^{k-1} \sin \pi h}{-h}$
 $= \lim_{h \rightarrow 0} \frac{(k-1)(-1)^{k-1} \sin \pi h}{-\pi h} \times \pi$
 $= -(k-1) (-1)^{k-1} \pi$
 $= (k-1) (-1)^k \pi$


$\because n$ is even
 $\sin(n\pi - \theta) = -\sin \theta$
 if n is odd
 $\sin(n\pi - \theta) = \sin \theta$

14.[D] $f(x) = \begin{cases} \sin x - x, & x \geq 0 \\ -\sin x + x, & x < 0 \end{cases}$

$f'(x) = \begin{cases} \cos x - 1, & x > 0 \\ -\cos x + 1, & x < 0 \end{cases}$
 $f'(0+) = 1 - 1 = 0$
 $f'(0-) = -1 + 1 = 0$
 \Rightarrow diff. at $x = 0$

15.[A] $f(x) = ||x| - 1|$

using graphical transformation



$\therefore f(x)$ is not differentiable at $x = 0, \pm 1$

$$16.[C] \quad g(x) = \frac{(x-1)^n}{\log \cos^m(x-1)}$$

Left hand derivative of $|x-1|$ at $x=1$ is $-1 = p$ (given)

$$\therefore \lim_{x \rightarrow 1^+} g(x) = p$$

$$\Rightarrow \lim_{h \rightarrow 0} \frac{h^n}{\log \cos^m h} = -1$$

$$\Rightarrow \lim_{h \rightarrow 0} \frac{nh^{n-1}}{m \cdot \frac{1}{\cosh} (-\sinh)} = -1 \quad (\text{applying D})$$

$$= \lim_{h \rightarrow 0} \frac{n}{m} \cdot \frac{h^{n-1} \cosh}{\sinh} = 1$$

If $n=2$ then

$$\lim_{h \rightarrow 0} \frac{2}{m} \cdot \frac{h}{\sinh} \cosh = 1$$

$$\Rightarrow \frac{2}{m} = 1$$

$$\Rightarrow m = 2$$

17.[B,C] $f: \mathbb{R} \rightarrow \mathbb{R}$

$$f(x+y) = f(x) + f(y)$$

$$\Rightarrow f(x) = \lambda x$$

Which of equation of straight line

Which is continuous & differentiable every where & $f'(x) = \lambda$ (constant function)

18.[A, B, C, D]

$$f(x) = \begin{cases} -x - \frac{\pi}{2}, & x \leq -\frac{\pi}{2} \\ -\cos x, & -\frac{\pi}{2} < x \leq 0 \\ x-1, & 0 < x \leq 1 \\ \ln x, & x > 1 \end{cases}$$

Option (A)

$$\text{at } x = -\frac{\pi}{2} \quad \text{L.H.L.} \quad \lim_{h \rightarrow 0} -\left(-\frac{\pi}{2} - h\right) - \frac{\pi}{2} = 0$$

$$\text{R.H.L.} \quad \lim_{h \rightarrow 0} -\cos\left(-\frac{\pi}{2} - h\right) = 0$$

$$f\left(-\frac{\pi}{2}\right) = -\left(-\frac{\pi}{2}\right) - \frac{\pi}{2} = 0$$

$$\therefore f(x) \text{ is continuous at } x = -\frac{\pi}{2}$$

Option (B)

LHD at $x=0$ is zero

R.H.D. at $x=0$ is 1

\therefore not diff. at $x=0$

Option (C)

L.H.D. at $x=1$ is 1

R.H.D. at $x=1$ is 1

differentiable at $x=1$

Option (D)

$$f'\left(-\frac{3}{2}\right) = \sin\left(-\frac{3}{2}\right) \text{ differentiable}$$

19.[B, D] At $x=2n$

$$x \rightarrow 2n^+ a_n + \sin 2n\pi = a_n$$

$$x \rightarrow 2n^- b_n + \cos 2n\pi = b_n + 1$$

For continuous $a_n = b_n + 1$

At $x=2n+1$

$$x \rightarrow 2n+1^+ b_{n+1} + \cos\pi(2n+1) = b_{n+1} - 1$$

$$x \rightarrow 2n+1^- a_n + \sin \pi(2n+1) = a_n$$

for continuous $a_n = b_{n+1} - 1$

$$a_n - b_{n+1} = -1$$

$$\text{for } n = n-1 \quad a_{n-1} - b_n = -1$$

$$20.[B] \quad f'(0+h) = \lim_{h \rightarrow 0} \frac{h^2 \left| \cos \frac{\pi}{h} \right| - 0}{h-0} = 0$$

$$f'(0-h) = \lim_{h \rightarrow 0} \frac{h^2 \left| \cos \frac{\pi}{h} \right| - 0}{-h} = 0$$

$$\therefore f'(0^+) = f'(0^-) = 0 = \text{finite}$$

So $f(x)$ is differentiable at $x=0$

$$f'(2+h) = \lim_{h \rightarrow 0} \frac{(2+h)^2 \left| \cos \left(\frac{\pi}{2+h} \right) \right| - 0}{h} = \pi$$

$$f'(2-h) = \lim_{h \rightarrow 0} \frac{(2-h)^2 \left| \cos \left(\frac{\pi}{2-h} \right) \right| - 0}{-h} = -\pi$$

$\therefore f'(2^+) \neq f'(2^-)$ but both are finite so $f(x)$ is not differentiable at $x=2$ but continuous at $x=2$