

INDEFINITE INTEGRATION

(KEY CONCEPTS + SOLVED EXAMPLES)

INDEFINITE INTEGRATION

1. Integration of a Function

2. Basic Theorems on Integration

3. Standard Integrals

4. Methods of Integration

5. Integrates of Different Expression

KEY CONCEPTS

1. Integration of a Function

Integration is a reverse process of differentiation. The **integral or primitive** of a function $f(x)$ with respect to x is that function (x) whose derivative with respect to x is the given function $f(x)$. It is expressed symbolically as -

$$\int f(x) dx = \phi(x)$$

Thus

$$\int f(x) dx = \phi(x) \Leftrightarrow \frac{d}{dx} [\phi(x)] = f(x)$$

The process of finding the **integral** of a function is called Integration and the given function is called **Integrand**. Now, it is obvious that the operation of integration is inverse operation of differentiation. Hence integral of a function is also named **anti-derivative** of that function.

Further we observe that-

$$\left. \begin{array}{l} \frac{d}{dx}(x^2) = 2x \\ \frac{d}{dx}(x^2 + 2) = 2x \\ \frac{d}{dx}(x^2 + k) = 2x \end{array} \right\} \Rightarrow \int 2x dx = x^2 + \text{constant}$$

So we always add a constant to the integral of function, which is called the **constant of Integration**. It is generally denoted by c . Due to presence of this constant such an integral is called an **Indefinite integral**.

2. Basic Theorems on Integration

If $f(x)$, $g(x)$ are two functions of a variable x and k is a constant, then-

$$(i) \quad \int k \cdot f(x) dx = k \int f(x) dx.$$

$$(ii) \quad \int [f(x) \pm g(x)] dx = \int f(x) dx \pm \int g(x) dx$$

$$(iii) \quad d/dx(\int f(x) dx) = f(x)$$

$$(iv) \quad \int \left(\frac{d}{dx} f(x) \right) dx = f(x)$$

3. Standard Integrals

The following integrals are directly obtained from the derivatives of standard functions.

$$i. \quad \int 0 \cdot dx = c$$

$$ii. \quad \int 1 \cdot dx = x + c$$

$$iii. \quad \int k \cdot dx = kx + c \quad (k \in R)$$

$$iv. \quad \int x^n dx = \frac{x^{n+1}}{n+1} + c \quad (n \neq -1)$$

$$v. \quad \int \frac{1}{x} dx = \log_e x + c$$

$$\text{vi. } \int e^x dx = e^x + c$$

$$\text{vii. } \int a^x dx = \frac{a^x}{\log_e a} + c = a^x \log_a e + c$$

$$\text{viii. } \int \sin x dx = -\cos x + c$$

$$\text{ix. } \int \cos x dx = \sin x + c$$

$$\text{x. } \int \tan x dx = \log \sec x + c = -\log \cos x + c$$

$$\text{xi. } \int \cot x dx = \log \sin x + c$$

$$\text{xii. } \begin{aligned} \int \sec x dx &= \log(\sec x + \tan x) + c \\ &= -\log(\sec x - \tan x) + c \\ &= \log \tan \left(\frac{\pi}{4} + \frac{x}{2} \right) + c \end{aligned}$$

$$\text{xiii. } \begin{aligned} \int \operatorname{cosec} x dx &= -\log (\operatorname{cosec} x + \cot x) + c \\ &= \log (\operatorname{cosec} x - \cot x) + c = \log \tan \left(\frac{x}{2} \right) + c \end{aligned}$$

$$\text{xiv. } \int \sec x \tan x dx = \sec x + c$$

$$\text{xv. } \int \operatorname{cosec} x \cot x dx = -\operatorname{cosec} x + c$$

$$\text{xvi. } \int \sec^2 x dx = \tan x + c$$

$$\text{xvii. } \int \operatorname{cosec}^2 x dx = -\cot x + c$$

$$\text{xviii. } \int \sinh x dx = \cosh x + c$$

$$\text{xix. } \int \cosh x dx = \sinh x + c$$

$$\text{xx. } \int \operatorname{sech}^2 x dx = \tanh x + c$$

$$\text{xxi. } \int \operatorname{cosech}^2 x dx = -\coth x + c$$

$$\text{xxii. } \int \operatorname{sech} x \tanh x dx = -\operatorname{sech} x + c$$

$$\text{xxiii. } \int \operatorname{cosech} x \coth x dx = -\operatorname{cosech} x + c$$

$$\text{xxiv. } \int \frac{1}{x^2 + a^2} dx = \frac{1}{a} \tan^{-1} \left(\frac{x}{a} \right) + c$$

$$\text{xxv. } \int \frac{1}{x^2 - a^2} dx = \frac{1}{2a} \log \left(\frac{x-a}{x+a} \right) + c$$

$$\text{xxvi. } \int \frac{1}{a^2 - x^2} dx = \frac{1}{2a} \log \left(\frac{x+a}{x-a} \right) + c$$

$$\text{xxvii. } \int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \left(\frac{x}{a} \right) + c$$

$$= -\cos^{-1} \left(\frac{x}{a} \right) + c$$

$$\text{xxviii. } \int \frac{1}{\sqrt{x^2 + a^2}} dx = \sinh^{-1} \left(\frac{x}{a} \right) + c$$

$$= \log(x + \sqrt{x^2 + a^2}) + c$$

$$\text{xxix. } \int \frac{1}{\sqrt{x^2 - a^2}} dx = \cosh^{-1} \left(\frac{x}{a} \right) + c$$

$$= \log(x + \sqrt{x^2 - a^2}) + c$$

$$\text{xxx. } \int \sqrt{a^2 - x^2} dx$$

$$= \frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \cdot \sin^{-1} \frac{x}{a} + c$$

$$\text{xxxi. } \int \sqrt{x^2 + a^2} dx$$

$$= \frac{x}{2} \sqrt{x^2 + a^2} + \frac{a^2}{2} \cdot \sinh^{-1} \frac{x}{a} + c$$

$$\text{xxxii. } \int \sqrt{x^2 - a^2} dx$$

$$= \frac{x}{2} \sqrt{x^2 - a^2} - \frac{a^2}{2} \cdot \cosh^{-1} \frac{x}{a} + c$$

$$\text{xxxiii. } \int \frac{1}{x \sqrt{x^2 - a^2}} dx = \frac{1}{a} \sec^{-1} \frac{x}{a} + c$$

$$\text{xxxiv. } \int e^{ax} \sin bx dx$$

$$= \frac{e^{ax}}{a^2 + b^2} (a \sin bx - b \cos bx) + c$$

$$= \frac{e^{ax}}{\sqrt{a^2 + b^2}} \sin \left\{ bx - \tan^{-1} \left(\frac{b}{a} \right) \right\} + c$$

$$\text{xxxv. } \int e^{ax} \cos bx dx$$

$$= \frac{e^{ax}}{a^2 + b^2} (a \cos bx + b \sin bx) + c$$

$$= \frac{e^{ax}}{\sqrt{a^2 + b^2}} \cos \left\{ bx - \tan^{-1} \frac{b}{a} \right\} + c$$

4. Methods of Integration

When integration can not be reduced into some standard form then integration is performed using following methods-

- (i) **Integration by substitutions**
- (ii) **Integration by parts**
- (iii) **Integration of rational functions**

- (iv) Integration of irrational functions
(v) Integration of trigonometric functions

4.1 INTEGRATION BY SUBSTITUTION:

Generally we apply this method in the following two cases.

- (i) When Integrand is a function of function -

$$\text{i.e. } \int f[\phi(x)] \phi'(x) dx$$

Here we put $\phi(x) = t$ so that the integrand $\phi'(x)$ is reduced to $\int f(t) dt$.

Note:

In this method the integrand is broken into two factors so that one factor can be expressed in terms of the function whose differential coefficient is the second factor.

- (ii) When integrand is the product of two factors such that one is the derivative of the other i.e.

$$I = \int f'(x)f(x) dx.$$

In this case we put $f(x) = t$ and convert it into a standard integral.

- (iii) Integral of a function of the form $f(ax + b)$.

Here we put $ax + b$ into standard integral. and convert it if
 $\int f(x) dx = \phi(x)$, then

$$\int f(ax+b) dx = \frac{1}{a} \phi(ax+b)$$

- (iv) Standard form of Integrals:

$$(a) \int \frac{f'(x)}{f(x)} dx = \log [f(x)] + c$$

$$(b) \int [f(x)]^n f'(x) dx = \frac{[f(x)]^{n+1}}{n+1} + c$$

(provided $n \neq -1$)

$$(c) \int \frac{f'(x)}{\sqrt{f(x)}} dx = 2\sqrt{f(x)} + c$$

- (v) Integral of the form

$$\int \frac{dx}{a \sin x + b \cos x}$$

putting $a = r \cos \theta$ and $b = r \sin \theta$. we get

$$I = \int \frac{dx}{r \sin(x+\theta)} = \frac{1}{r} \int \cosec(x+\theta) dx .$$

$$= \frac{1}{r} \log \tan(x/2 + \theta/2) + c$$

$$= \frac{1}{\sqrt{a^2+b^2}} \log \tan(x/2 + 1/2 \tan^{-1} b/a) + c$$

- (vi) Standard Substitutions : Following standard substitutions will be useful-

Integrand form	Substitution
(i) $\sqrt{a^2 - x^2}$ or	$x = a \sin \theta$ or

- $\frac{1}{\sqrt{a^2 - x^2}}$ $x = a \cos \theta$
(ii) $\sqrt{x^2 + a^2}$ or $x = a \tan \theta$ or
 $\frac{1}{\sqrt{x^2 + a^2}}$ $x = a \cot \theta$ or $x = a \sinh \theta$
(iii) $\sqrt{x^2 - a^2}$ or $x = a \sec \theta$ or
 $\frac{1}{\sqrt{x^2 - a^2}}$ $x = \operatorname{acosec} \theta$ or $x = a \cosh \theta$
(iv) $\sqrt{\frac{x}{a+x}}$ or $\sqrt{\frac{a+x}{x}}$ $x = a \tan^2 \theta$
or $\sqrt{x(a+x)}$ or $\frac{1}{\sqrt{x(a+x)}}$
(v) $\sqrt{\frac{x}{a-x}}$ or $\sqrt{\frac{a-x}{x}}$
or $\sqrt{x(a-x)}$ $x = a \sin^2 \theta$
or $\frac{1}{\sqrt{x(a-x)}}$
(vi) $\sqrt{\frac{x}{x-a}}$ or $x = a \sec^2 \theta$
 $\sqrt{\frac{x-a}{x}}$ or
 $\sqrt{x(x-a)}$ or
(vii) $\sqrt{\frac{a-x}{a+x}}$ or $x = a \cos 2\theta$
 $\sqrt{\frac{a+x}{a-x}}$
(viii) $\sqrt{\frac{x-\alpha}{\beta-x}}$ or $x = \alpha \cos^2 \theta + \beta \sin^2 \theta$
 $\sqrt{(\alpha-x)(\beta-x)}$
 $(\beta > \alpha)$

4.2 INTEGRATION BY PARTS :

4.2.1 If u and v are two functions of x , then

$$\int (u.v) dx = u \left(\int v dx \right) - \int \left(\frac{du}{dx} \right) \cdot \left(\int v dx \right) dx.$$

i.e. Integral of the product of two functions
= first function \times integral of second function
 $- \int [(\text{derivative of first}) \times (\text{Integral of second})]$

Note :

- (i) From the first letter of the words inverse circular, logarithmic, Algebraic, Trigonometric, Exponential functions, we get a word **ILATE**. Therefore first arrange the functions in the order according to letters of this word and then integrate by parts.
- (ii) For the integration of Logarithmic or Inverse trigonometric functions alone, take unity (1) as the second function.

4.2.2 If the integral is of the form $\int e^x [f(x) + f'(x)]dx$, then by breaking this integral into two integrals, integrate one integral by parts and keep other integral as it is, By doing so, we get-

$$\boxed{\int e^x [f(x) + f'(x)]dx = e^x f(x) + c}$$

4.2.3 If the integral is of the form $\int [xf'(x) + f(x)]dx$ then by breaking this integral into two integrals integrate one integral by parts and keep other integral as it is, by doing so, we get

$$\boxed{\int [xf'(x) + f(x)]dx = x f(x) + c}$$

4.3 Integration of Rational functions:

4.3.1 When denominator can be factorized (using partial fraction) :

Let the integrand is of the form $\frac{f(x)}{g(x)}$, where both $f(x)$ and $g(x)$ are polynomials. If degree of $f(x)$ is greater than degree of $g(x)$ then first divide $f(x)$ by $g(x)$ till the degree of the remainder becomes less than the degree of $g(x)$. Let $Q(x)$ is the quotient and $R(x)$, the remainder then

$$\frac{f(x)}{g(x)} = Q(x) + \frac{R(x)}{g(x)}$$

Now in $R(x)/g(x)$, factorize $g(x)$ and then write partial fractions in the following manner-

(i) For every non repeated linear factor in the denominator, write

$$\frac{1}{(x-a)(x-b)} = \frac{A}{x-a} + \frac{B}{x-b}$$

(ii) For repeated linear factors in the denominator, write-

$$\frac{1}{(x-a)^3(x-b)} = \frac{A}{(x-a)} + \frac{B}{(x-a)^2}$$

$$+ \frac{C}{(x-a)^3} + \frac{D}{(x-b)}$$

(iii) For every non repeated quadratic factor in the denominator, write

$$\frac{1}{(ax^2 + bx + c)(x-d)} = \frac{Ax + B}{ax^2 + bx + c} + \frac{C}{x-d}$$

Note :

(i) If integrand is of the form $\frac{1}{(x+a)(x+b)}$ then use the following method for obtaining partial fractions-

$$\text{Here } \frac{1}{(x+a)(x+b)} = \frac{1}{(a-b)} \left[\frac{a-b}{(x+a)(x+b)} \right]$$

$$= \frac{1}{(a-b)} \left[\frac{(x+a)-(x+b)}{(x+a)(x+b)} \right]$$

$$= \frac{1}{(a-b)} \left[\frac{1}{x+b} - \frac{1}{x+a} \right]$$

(ii) If integrand is of the form $\frac{x}{(x+a)(x+b)}$ then

$$\frac{x}{(x+a)(x+b)} = \frac{1}{b-a} \left[\frac{(b-a)x}{(x+a)(x+b)} \right]$$

$$\begin{aligned}
 &= \frac{1}{b-a} \left[\frac{b(x+a) - a(x+b)}{(x+a)(x+b)} \right] \\
 &= \frac{1}{b-a} \left[\frac{b}{x+b} - \frac{a}{x+a} \right]
 \end{aligned}$$

4.3.2 When denominator can not be factorised:

In this case integral may be in the form

$$(i) \int \frac{dx}{ax^2 + bx + c}, (ii) \int \frac{(px+q)}{ax^2 + bx + c} dx$$

Method:

- (i) Here taking coefficient of x^2 common from denominator, write -

$$x^2 + (b/a)x + c/a = (x + b/2a)^2 - \frac{b^2 - 4ac}{4a^2}$$

Now the integrand so obtained can be evaluated easily by using standard formulas.

- (ii) Here suppose that $px + q = A$ [diff. coefficient of $(ax^2 + bx + c)$] + B

$$= A(2ax + b) + B \quad \dots(1)$$

Now comparing coefficient of x and constant terms.

we get $A = p/2a$, $B = q - (pb/2a)$

$$\begin{aligned}
 \therefore I &= P/2a \int \frac{2ax + b}{ax^2 + bx + c} dx \\
 &\quad + \left(q - \frac{pb}{2a} \right) \int \frac{dx}{ax^2 + bx + c}
 \end{aligned}$$

Now we can integrate it easily.

4.3.3 Integration of rational functions containing only even powers of x.

To find integral of such functions, first we divide numerator and denominator by x^2 , then express numerator as $d(x \pm 1/x)$ and denominator as a function of $(x \pm 1/x)$.

4.4 Integration of irrational functions :

If anyone term in Nr or Dr is irrational then it is made rational by suitable substitution. Also if integral is of the form-

$$\int \frac{dx}{\sqrt{ax^2 + bx + c}}, \int \sqrt{ax^2 + bx + c} dx$$

then we integrate it by expressing

$$ax^2 + bx + c = (x + \alpha)^2 + \beta$$

Also for integrals of the form

$$\int \frac{px+q}{\sqrt{ax^2 + bx + c}} dx, \int (px+q) \sqrt{ax^2 + bx + c} dx.$$

First we express $px + q$ in the form

$$px + q = A \left\{ \frac{d}{dx}(ax^2 + bx + c) \right\} + B \text{ and then proceed as usual with standard form.}$$

4.5 Integration of Trigonometric functions :

Here we shall study the methods for evaluation of following types of integrals.

I. (i) $\int \frac{dx}{a + b \sin^2 x}$

$$(ii) \int \frac{dx}{a + b \cos^2 x}$$

$$(iii) \int \frac{dx}{a \cos^2 x + b \sin x \cos x + c \sin^2 x}$$

$$(iv) \int \frac{dx}{(a \sin x + b \cos x)^2}$$

Method :

Divide numerator and Denominator by $\cos^2 x$ in all such type of integrals and then put $\tan x = t$.

$$II. (i) \int \frac{dx}{a + b \cos x}$$

$$(ii) \int \frac{dx}{a + b \sin x}$$

$$(iii) \int \frac{dx}{a \cos x + b \sin x}$$

$$(iv) \int \frac{dx}{a \sin x + b \cos x + c}$$

Method :

In such types of integrals we use following formulae for $\sin x$ and $\cos x$ in terms of $\tan(x/2)$.

$$\sin x = \frac{2 \tan\left(\frac{x}{2}\right)}{1 + \tan^2\left(\frac{x}{2}\right)}, \cos x = \frac{1 - \tan^2\left(\frac{x}{2}\right)}{1 + \tan^2\left(\frac{x}{2}\right)}$$

and then take $\tan(x/2) = t$ and integrate another method for evaluation of integral

(iii) put $a = r \cos \alpha$, $b = r \sin \alpha$, then

$$\begin{aligned} I &= \frac{1}{r} \int \frac{dx}{\sin(x + \alpha)} \\ &= \frac{1}{r} \int \cosec(x + \alpha) dx . \\ &= \frac{1}{r} \log \tan(x/2 + \alpha/2) + C \\ &= \frac{1}{\sqrt{a^2 + b^2}} \log \tan\left(\frac{x}{2} + \frac{1}{2} \tan^{-1} \frac{b}{a}\right) + C \end{aligned}$$

$$III. \int \frac{p \sin x + q \cos x}{a \sin x + b \cos x} dx$$

$$\int \frac{p \sin x}{a \sin x + b \cos x} dx$$

$$\int \frac{q \cos x}{a \sin x + b \cos x} dx$$

For their integration, we first express Nr. as follows-

Nr = A (Dr) + B (derivative of Dr.)

Then integral = Ax + B log (Dr) + C

5. Some Integrates of Different Expression of e^x

$$(i) \int \frac{ae^x}{b + ce^x} dx \quad [\text{put } e^x = t]$$

- (ii) $\int \frac{1}{1+e^x} dx$ [Multiply and divide by e^{-x} and put $e^{-x}=t$]
- (iii) $\int \frac{1}{1-e^x} dx$ [Multiply and divide by e^{-x} and put $e^{-x}=t$]
- (iv) $\int \frac{1}{e^x - e^{-x}} dx$ [Multiply and divide by e^x]
- (v) $\int \frac{e^x - e^{-x}}{e^x + e^{-x}} dx$ $\left[\frac{f'(x)}{f(x)} \text{ form} \right]$
- (vi) $\int \frac{e^x + 1}{e^x - 1} dx$ [Multiply and divide by $e^{-x/2}$]
- (vii) $\int \left(\frac{e^x - e^{-x}}{e^x + e^{-x}} \right)^2 dx$ [Integrand = $\tanh^2 x$]
- (viii) $\int \left(\frac{e^{2x} + 1}{e^{2x} - 1} \right)^2 dx$ [Integrand = $\coth^2 x$]
- (ix) $\int \frac{1}{(e^x + e^{-x})^2} dx$ [Integrand = $1/4 \operatorname{sech}^2 x$]
- (x) $\int \frac{1}{(e^x - e^{-x})^2} dx$ [Integrand = $1/4 \operatorname{cosech}^2 x$]
- (xi) $\int \frac{1}{(1+e^x)(1-e^{-x})} dx$ [Multiply and divide by e^x
and put $e^x = t$]
- (xii) $\int \frac{1}{\sqrt{1-e^x}} dx$ [Multiply and divide by $e^{-x/2}$]
- (xiii) $\int \frac{1}{\sqrt{1+e^x}} dx$ [Multiply and divide by $e^{-x/2}$]
- (xiv) $\int \frac{1}{\sqrt{e^x - 1}} dx$ [Multiply & divide by $e^{-x/2}$]
- (xv) $\int \frac{1}{\sqrt{2e^x - 1}} dx$ [Multiply and divide by $\sqrt{2} e^{-x/2}$]
- (xvi) $\int \sqrt{1-e^x} dx$ [Integrand
 $= (1 - e^x)/\sqrt{1-e^x}$]
- (xvii) $\int \sqrt{1+e^x} dx$ [Integrand
 $= (1 + e^x)/\sqrt{1+e^x}$]
- (xviii) $\int \sqrt{e^x - 1} dx$ [Integrand
 $= (e^x - 1)/\sqrt{e^x - 1}$]

$$(xix) \int \sqrt{\frac{e^x + a}{e^x - a}} dx \quad [\text{Integrand}]$$
$$= (e^x + a) / \sqrt{e^{2x} - a^2}$$

SOLVED EXAMPLE

Ex.1 $\int \sin^2(x/2) dx$ equals-

- (A) $\frac{1}{2}(x + \sin x) + c$ (B) $\frac{1}{2}(x + \cos x) + c$
 (C) $\frac{1}{2}(x - \sin x) + c$ (D) None of these

Sol. Here $I = \int \frac{1-\cos x}{2} dx$

$$= \frac{1}{2}(x - \sin x) + c$$

Ans.[C]

Ex.2 $\int \cot^2 x dx$ equals -

- (A) $-\sec x + x + c$ (B) $-\cot x - x + c$
 (C) $-\sin x + x + c$ (D) None of these

Sol. $\int (\operatorname{cosec}^2 x - 1) dx$

$$= -\cot x - x + c$$

Ans. [B]

Ex.3 $\int \frac{5x+7}{x} dx$ equals-

- (A) $5x + 7 \log x$ (B) $7x + 5 \log x + c$
 (C) $5x + 7 \log x + c$ (D) None of these

Sol. $\int \frac{5x+7}{x} dx = \int \left(\frac{5x}{x} + \frac{7}{x} \right) dx$

$$= \int 5 dx + \int \frac{7}{x} dx = 5 \int 1 dx + 7 \int \frac{1}{x} dx$$

$$= 5x + 7 \log x + c$$

Ans.[C]

Ex.4 $\int \left(x - \frac{1}{x} \right)^3 dx$, ($x > 0$) equals-

- (A) $\frac{x^3}{3} - \frac{3}{2}x^2 + 3 \log x + \frac{1}{2x^2} + c$
 (B) $\frac{x^4}{3} - \frac{3}{2}x^2 + 3 \log x + \frac{1}{2x^2} + c$
 (C) $\frac{x^4}{4} + 3 \log x + \frac{1}{2x^2} + c$
 (D) None of these

Sol. $\int \left(x - \frac{1}{x} \right)^3 dx$
 $= \int \left(x^3 - 3x^2 \cdot \frac{1}{x} + 3x \cdot \frac{1}{x^2} - \frac{1}{x^3} \right) dx$
 $[\because (a-b)^3 = (a^3 - 3a^2b + 3ab^2 - b^3)]$

$$\begin{aligned} &= \int \left(x^3 - 3x + \frac{3}{x} - \frac{1}{x^3} \right) dx \\ &= \int x^3 dx - 3 \int x dx + 3 \int \frac{1}{x} dx - \int \frac{1}{x^3} dx \\ &= \frac{x^{3+1}}{3+1} - 3 \cdot \frac{x^{1+1}}{1+1} + 3 \log x - \frac{x^{-3+1}}{-3+1} + c \\ &= \frac{x^4}{4} - \frac{3}{2}x^2 + 3 \log x + \frac{1}{2x^2} + c \end{aligned}$$

Ans.[B]

Ex.5 The value of $\int \left(\frac{6}{1+x^2} + 10^x \right) dx$ is -

(A) $6 \tan^{-1} x + 10^x \log_e 10 + c$

(B) $6 \tan^{-1} x + \frac{10^x}{\log_e 10} + c$

(C) $3 \tan^{-1} x + \frac{10^x}{\log_e 10} + c$

(D) None of these

Sol. $\int \left(\frac{6}{1+x^2} + 10^x \right) dx$
 $= 6 \int \frac{1}{1+x^2} dx + \int 10^x dx$
 $= 6 \tan^{-1} x + \frac{10^x}{\log_e 10} + C$

Ans.[B]

Ex.6 $\int (\tan x + \cot x)^2 dx$ is equal to-

- (A) $\tan x - \cot x + c$ (B) $\tan x + \cot x + c$
 (C) $\cot x - \tan x + c$ (D) None of these

Sol. $I = \int (\tan^2 x + \cot^2 x + 2) dx$

$$= \int (\sec^2 x + \operatorname{cosec}^2 x) dx$$



$$= \tan x - \cot x + c$$

Ans. [A]

Sol. $\int \sec^2(ax + b) dx$, putting $ax + b = t$,

$$adx + 0 = dt \text{ or } dx = \frac{dt}{a}$$

$$\therefore \int \sec^2(ax + b) dx = \int \sec^2 t \frac{dt}{a} \\ = \frac{1}{a} \int \sec^2 t dt$$

$$= \frac{1}{a} \tan t + c$$

$$= \frac{1}{a} \tan(ax + b) + c$$

(Putting the value of t)

Ans.[C]

Ex.7 $\int \sin 2x \sin 3x dx$ equals-

$$(A) \frac{1}{2} (\sin x - \sin 5x) + c$$

$$(B) \frac{1}{10} (\sin x - \sin 5x) + c$$

$$(C) \frac{1}{10} (5 \sin x - \sin 5x) + c$$

(D) None of these

$$\text{Sol. } I = \frac{1}{2} \int [\cos(-x) - \cos 5x] dx$$

$$= \frac{1}{2} \left[\sin x - \frac{\sin 5x}{5} \right] + c$$

$$= \frac{1}{10} [5 \sin x - \sin 5x] + c$$

Ans. [C]

Ex.8 $\int \frac{x^2}{x^2 - 1} dx$ equals-

$$(A) x + \log \sqrt{\frac{x-1}{x+1}} + c \quad (B) x + \log \sqrt{\frac{x+1}{x-1}} + c$$

$$(C) x + \log \left(\frac{x-1}{x+1} \right) + c \quad (D) x + \log \left(\frac{x+1}{x-1} \right) + c$$

$$\text{Sol. } \int \frac{x^2 - 1 + 1}{x^2 - 1} dx$$

$$= \int \left(1 + \frac{1}{x^2 - 1} \right) dx$$

$$= x + \frac{1}{2} \log \left(\frac{x-1}{x+1} \right) + c$$

$$= x + \log \sqrt{\frac{x-1}{x+1}} + c$$

Ans.[A]

Ex.9 $\int \sec^2(ax + b) dx$ equals-

$$(A) \tan(ax + b) + c \quad (B) \frac{1}{2} \tan x + c$$

$$(C) \frac{1}{a} \tan(ax + b) + c \quad (D) \text{None of these}$$

Ex.10 $\int \frac{1}{x \log x} dx$ is equal to-

$$(A) \log(x \log x) + c \quad (B) \log(\log x + x) + c$$

$$(C) \log x + c \quad (D) \log(\log x) + c$$

$$\text{Sol. } \int \frac{1}{x \log x} dx = \int \frac{1}{x} \cdot \frac{1}{\log x} dx$$

$$\text{put } \log x = t, \frac{1}{x} dx = dt$$

$$\therefore \int \frac{1}{x} \cdot \frac{1}{\log x} dx = \int \frac{1}{t} dt$$

$$\therefore \int \frac{1}{t} dt = \log t + c = \log(\log x) + c$$

(putting the value of t = log x)

Ans.[D]

Ex.11 $\int \sec^2 x \cos(\tan x) dx$ equals-

$$(A) \sin(\cos x) + c \quad (B) \sin(\tan x) + c$$

$$(C) \operatorname{cosec}(\tan x) + c \quad (D) \text{None of these}$$

Sol. Let $\tan x = t$, then $\sec^2 x dx = dt$

$$\therefore I = \int \cos t dt = \sin t + c$$

$$= \sin(\tan x) + c$$

Ans.[B]

Ex.12 $\int \tan^n x \sec^2 x dx$ equals-

$$(A) \frac{\tan^{n-1} x}{n-1} + c \quad (B) \frac{\tan^{n-1} x}{n+1} + c$$



- Sol.** (C) $\tan^{n+1} x + c$ (D) None of these
 $\int \tan^n x \sec^2 x dx$

putting $\tan x = t$, $\sec^2 x dx = dt$

$$\begin{aligned}\int \tan^n x \sec^2 x dx &= \int t^n dt = \frac{\tan^{n+1}}{n+1} + c \\ &= \frac{(\tan x)^{n+1}}{n+1} + c\end{aligned}$$

Ans.[B]

- Ex.13** $\int \frac{\sin 2x}{1+\cos^4 x} dx$ is equal to-

- (A) $\cos^{-1}(\cos^2 x) + c$ (B) $\sin^{-1}(\cos^2 x) + c$
(C) $\cot^{-1}(\cos^2 x) + c$ (D) None of these

- Sol.** Here differential coefficient of $\cos^2 x$ is $-\sin 2x$

Let $\cos^2 x = t$

$$\therefore 2 \cos x (-\sin x) dx = dt$$

$$\text{or } \sin 2x dx = -dt$$

$$\begin{aligned}\therefore \int \frac{\sin 2x}{1+\cos^4 x} dx &= \int \frac{-dt}{1+t^2} \\ &= \cot^{-1} t + c \\ &= \cot^{-1} (\cos^2 x) + c\end{aligned}$$

Ans.[C]

- Ex.14** $\int \frac{be^x}{\sqrt{a+be^x}} dx$ equals-

- (A) $\frac{2}{b} \sqrt{a+be^x} + c$ (B) $\frac{1}{b} \cdot \sqrt{a+be^x} + c$
(C) $2 \sqrt{a+be^x} + c$ (D) None of these

- Sol.** $\int \frac{be^x}{\sqrt{a+be^x}} dx$, putting $a+be^x=t$

$$be^x dx = dt$$

$$\begin{aligned}\therefore \int \frac{be^x}{\sqrt{a+be^x}} dx &= \int \frac{dt}{\sqrt{t}} = 2\sqrt{t} + c \\ &= 2\sqrt{a+be^x} + c\end{aligned}$$

Ans.[C]

- Ex.15** $\int \sqrt{\frac{1+\cos x}{1-\cos x}} dx$ equals-

- (A) $\log \cos\left(\frac{x}{2}\right) + c$ (B) $2\log \sin\left(\frac{x}{2}\right) + c$
(C) $2 \log \sec\left(\frac{x}{2}\right) + c$ (D) None of these

$$\begin{aligned}\text{Sol. } I &= \int \sqrt{\frac{1+\cos x}{1-\cos x}} dx \\ &= \int \sqrt{\frac{2\cos^2(x/2)}{2\sin^2(x/2)}} dx \\ &= \int \cot\left(\frac{x}{2}\right) dx \\ &= 2 \log \sin\left(\frac{x}{2}\right) + c\end{aligned}$$

Ans.[B]

- Ex.16** $\int \frac{\sqrt{\tan x}}{\sin x \cos x} dx$ equals-

- (A) $2\sqrt{\sec x} + c$ (B) $2\sqrt{\tan x} + c$
(C) $2/\sqrt{\tan x} + c$ (D) $2/\sqrt{\sec x} + c$

$$\begin{aligned}\text{Sol. } I &= \int \frac{\sqrt{\tan x}}{\tan x} \sec^2 x dx \\ &= \int \frac{\sec^2 x}{\sqrt{\tan x}} dx = 2\sqrt{\tan x} + c\end{aligned}$$

Ans.

[B]

- Ex.17** $\int \sin^5 x \cdot \cos^3 x dx$ is equal to-

- (A) $\frac{\sin^6 x}{6} - \frac{\sin^8 x}{8} + c$ (B) $\frac{\cos^6 x}{6} - \frac{\cos^8 x}{8} + c$
(C) $\frac{\cos^6 x}{6} - \frac{\sin^8 x}{8} + c$ (D) None of these

- Sol.** $\int \sin^5 x \cdot \cos^3 x dx$

Assumed that $\sin x = t$

$$\therefore \cos x dx = dt$$

$$\begin{aligned}&= \int t^5(1-t^2) dt = \int (t^5 - t^7) dt \\ &= \frac{t^6}{6} - \frac{t^8}{8} + c\end{aligned}$$

$$= \frac{\sin^6 x}{6} - \frac{\sin^8 x}{8} + c$$

Ans.[A]

Ex.18 $\int \frac{x^2}{1+x^6} dx$ is equal to-

- (A) $\tan^{-1}x^3 + c$ (B) $\tan^{-1}x^2 + c$

$$(C) \frac{1}{3} \tan^{-1}x^3 + c \quad (D) 3 \tan^{-1}x^3 + c$$

Sol. Put $x^3 = t \Rightarrow x^2 dx = \frac{1}{3} dt$

$$\therefore I = \frac{1}{3} \int \frac{dt}{1+t^2} = \frac{1}{3} \tan^{-1} x^3 + c$$

Ans.

[C]

Ex.19 $\int \sqrt{\frac{1+x}{1-x}} dx$ equals-

- (A) $\sin^{-1} x + \sqrt{1-x^2} + c$
 (B) $\sin^{-1} x + \sqrt{x^2-1} + c$
 (C) $\sin^{-1} x - \sqrt{1-x^2} + c$
 (D) $\sin^{-1} x - \sqrt{x^2-1} + c$

$$I = \int \sqrt{\frac{1+x}{1-x}} dx$$

$$= \int \frac{dx}{\sqrt{1-x^2}} - \frac{1}{2} \int \frac{-2x dx}{\sqrt{1-x^2}}$$

$$= \sin^{-1} x - \sqrt{1-x^2} + c$$

Ans.

[C]

Ex.20 The primitive of $\log x$ will be-

- (A) $x \log(e+x) + c$ (B) $x \log\left(\frac{e}{x}\right) + c$
 (C) $x \log\left(\frac{x}{e}\right) + c$ (D) $x \log(ex) + c$

$$\text{Sol. } \int \log x dx = \int \log x \cdot 1 dx$$

[Integrating by parts, taking $\log x$ as first part and 1 as second part]

$$= (\log x) \cdot x - \int \left\{ \frac{d(\log x)}{dx} \right\} \cdot x dx$$

$$= x \log x - \int \frac{1}{x} \cdot x dx = (x \log x - x) + c$$

$$= x(\log x - 1) + c = \log\left(\frac{x}{e}\right) + c$$

Ans. [C]

Ex.21 $\int x \tan^{-1} x$ is equal to-

$$(A) \frac{1}{2}(x^2 + 1) \tan^{-1} x - x + c$$

$$(B) \frac{1}{2}(x^2 + 1) \tan^{-1} x + x + c$$

$$(C) \frac{1}{2}(x^2 + 1) \tan^{-1} x - \frac{1}{2}x + c$$

$$(D) \frac{1}{2}(x^2 - 1) \tan^{-1} x - \frac{1}{2}x + c$$

Sol. Integrating by parts taking x as second part

$$I = \frac{x^2}{2} \tan^{-1} x - \int \frac{1}{1+x^2} \cdot \frac{x^2}{2} dx$$

$$= \frac{1}{2}x^2 \tan^{-1} x - \frac{1}{2} \left(1 - \frac{1}{1+x^2} \right) dx$$

$$= \frac{1}{2}x^2 \tan^{-1} x - \frac{1}{2}x + \frac{1}{2} \tan^{-1} x + c$$

$$= \frac{1}{2}(x^2 + 1) \tan^{-1} x - \frac{1}{2}x + c \quad \text{Ans. [C]}$$

Ex.22 $\int \sin(\log x) dx$ equals-

$$(A) \frac{x}{\sqrt{2}} \sin(\log x + \frac{\pi}{8}) + c$$

$$(B) \frac{x}{\sqrt{2}} \sin(\log x - \frac{\pi}{4}) + c$$

$$(C) \frac{x}{\sqrt{2}} \cos(\log x - \frac{\pi}{4}) + c$$

(D) None of these

Sol. $\int \sin(\log x) dx$, assumed that $x = e^t$

$$\therefore dx = e^t dt$$

$$= \int \sin t \cdot e^t dt$$

$$= \frac{e^t}{\sqrt{1+1}} \sin(t - \tan^{-1} 1) + c$$

$$\Rightarrow \int \sin(\log x) dx$$

$$= \frac{x}{\sqrt{2}} \sin \left(\log x - \frac{\pi}{4} \right) + c$$

Ans.

[B]

Ex.23 $\int \frac{xe^x}{(x+1)^2} dx$ is equal to-

(A) $\frac{e^x}{(x+1)^2} + c$ (B) $\frac{e^x}{x+1} + c$

(C) $\frac{e^x}{(x+1)^2} + c$ (D) None of these

Sol. $I = \int e^x \left[\frac{(x+1)-1}{(x+1)^2} \right] dx$

$$= \int e^x \left(\frac{1}{x+1} + \frac{-1}{(x+1)^2} \right) dx$$

$$= e^x f(x) + c$$

$$= \frac{e^x}{x+1} + c$$

Ans. [B]

Ex. 24 $\int x^3 (\log x)^2 dx$ equals-

(A) $\frac{1}{32} x^4 [8 (\log x)^2 - 4 \log x + 1] + c$

(B) $\frac{1}{32} x^4 [8 (\log x)^2 - 4 \log x - 1] + c$

(C) $\frac{1}{32} x^4 [8 (\log x)^2 + 4 \log x + 1] + c$

(D) None of these

Sol. Integrating by parts taking x^3 as second part

$$I = \frac{1}{4} x^4 (\log x)^2 - \frac{1}{2} \int x^3 \log x dx$$

$$= \frac{1}{4} x^4 (\log x)^2 - \frac{1}{2} \left(\frac{1}{4} x^4 \log x - \frac{1}{16} x^4 \right) + c$$

$$= \frac{1}{32} x^4 [8 (\log x)^2 - 4 \log x + 1] + c$$

Ans. [A]

Ex.25 The value of $\int x \sec x \tan x dx$ is-

(A) $x \sec x + \log (\sec x + \tan x) + c$

(B) $x \sec x - \log (\sec x - \tan x) + c$

(C) $x \sec x + \log (\sec x - \tan x) + c$

(D) None of the above

Sol. $\int x \cdot (\sec x \tan x) dx$

$$= (x \cdot \sec x) - \int (1 \cdot \sec x) dx$$

(Integrating by parts, taking x as first function)

$$= x \sec x - \log (\sec x + \tan x) + c$$

$$= x \sec x - \log \left\{ (\sec x + \tan x) \frac{\sec x - \tan x}{\sec x + \tan x} \right\} + c$$

$$= x \sec x - \log \left(\frac{\sec^2 x - \tan^2 x}{\sec x + \tan x} \right) + c$$

$$= x \sec x + \log (\sec x - \tan x) + c$$

Ans.

[C]

Ex.26 $\int \frac{x + \sin x}{1 + \cos x} dx$ equals-

(A) $\frac{x}{2} \tan \left(\frac{x}{2} \right) + c$ (B) $\frac{x}{2} \tan x + c$

(C) $x \tan \left(\frac{x}{2} \right) + c$ (D) $x \tan x + c$

Sol. $I = \int \frac{x + 2 \sin(x/2) \cos(x/2)}{2 \cos^2(x/2)} dx$

$$= \frac{1}{2} \int x \sec^2(x/2) dx + \int \tan(x/2) dx$$

$$= x \tan(x/2) - \int \tan(x/2) dx + \int \tan(x/2) dx$$

$$= x \tan \left(\frac{x}{2} \right) + c.$$

Ans.[C]

Ex.27 $\int e^x \frac{x-1}{(x+1)^3} dx$ equals-

(A) $-\frac{e^x}{x+1} + c$ (B) $\frac{e^x}{x+1} + c$

(C) $\frac{e^x}{(x+1)^2} + c$ (D) $-\frac{e^x}{(x+1)^2} + c$

Sol. $I = \int e^x \left[\frac{x+1-2}{(x+1)^3} \right] dx$

$$= \int e^x \left(\frac{1}{(x+1)^2} - \frac{2}{(x+1)^3} \right) dx$$

Thus the given integral is of the form

$$= \int e^x \{f(x) + f'(x)\} dx$$

$$\therefore I = e^x f(x) = \frac{e^x}{(x+1)^2} + c$$

Ans.[C]

Ex. 28 $\int \sec^3 \theta d\theta$ is equal to-

- (A) $\frac{1}{2} [\tan \theta \sec \theta + \log(\tan \theta + \sec \theta)] + c$
- (B) $\frac{1}{2} \tan \theta \sec \theta + \log(\tan \theta + \sec \theta) + c$
- (C) $\frac{1}{2} [\tan \theta \sec \theta - \log(\tan \theta + \sec \theta)] + c$
- (D) None of these

Sol. $I = \int \sec \theta \sec^2 \theta d\theta$

$$= \int \sqrt{\tan^2 \theta + 1} \sec^2 \theta d\theta$$

$$= \int \sqrt{t^2 + 1} dt, \text{ where } t = \tan \theta$$

$$= \frac{t}{2} \sqrt{t^2 + 1} + \frac{1}{2} \log(t + \sqrt{t^2 + 1}) + c$$

$$= \frac{1}{2} [\tan \theta \sec \theta + \log(\tan \theta + \sec \theta)] + c$$

Ans. [A]

Ex.29 $\int \frac{\cos x + x \sin x}{x(x + \cos x)} dx$ is equal to-

- (A) $\log \{x(x + \cos x)\} + c$
- (B) $\log \left(\frac{x}{x + \cos x} \right) + c$
- (C) $\log \left(\frac{x + \cos x}{x + \cos x} \right) + c$
- (D) None of these

Sol. $I = \int \frac{(x + \cos x) - x + x \sin x}{x(x + \cos x)} dx$

$$= \int \frac{1}{x} dx - \int \frac{1 - \sin x}{x + \cos x} dx$$

$$= \log x - \log(x + \cos x) + c$$

$$= \log \left(\frac{x}{x + \cos x} \right) + c$$

[B]

Ex.30 $\int \sqrt{\sec x - 1} dx$ is equal to-

- (A) $2 \sin^{-1}(\sqrt{2} \cos x / 2) + c$
- (B) $-2 \sinh^{-1}(\sqrt{2} \cos x / 2) + c$
- (C) $-2 \cosh^{-1}(\sqrt{2} \cos x / 2) + c$
- (D) None of these

Sol. $I = \int \sqrt{\frac{1 - \cos x}{\cos x}} dx$

$$= \int \frac{\sqrt{2} \sin x / 2}{\sqrt{2 \cos^2 x / 2 - 1}} dx$$

$$= -2 \int \frac{dt}{\sqrt{t^2 - 1}} \text{ where } t = \sqrt{2} \cos x / 2$$

$$= -2 \cosh^{-1} t + c$$

$$= -2 \cosh^{-1} (\sqrt{2} \cos x / 2) + c$$

Ans.

Ex.31 $\int \frac{x^2 + 1}{(x-1)(x-2)} dx$ equals-

- (A) $\log \left[\frac{(x-2)^5}{(x-1)^2} \right] + c$
- (B) $x + \log \left[\frac{(x-2)^5}{(x-1)^2} \right] + c$
- (C) $x + \log \left[\frac{(x-1)^5}{(x-2)^5} \right] + c$

- (D) None of these

Sol.

Here since the highest powers of x in Num^r and Den^r are equal and coefficients of x² are also equal, therefore

$$\frac{x^2 + 1}{(x-1)(x-2)} \equiv 1 + \frac{A}{x-1} + \frac{B}{x-2}$$

On solving we get A = -2, B = 5

$$\text{Thus } \frac{x^2 + 1}{(x-1)(x-2)} \equiv 1 - \frac{2}{x-1} + \frac{5}{x-2}$$

The above method is used to obtain the value of constant corresponding to non repeated linear factor in the Den^r.

Ans.

$$\text{Now } I = \left(1 - \frac{2}{x-1} + \frac{5}{x-2} \right) dx$$

$$= x - 2 \log(x-1) + 5 \log(x-2) + c$$

$$= x + \log \left[\frac{(x-2)^5}{(x-1)^2} \right] + c$$

Ans.[B]

Ex.32 The value of $\int \frac{x^2 dx}{(x^2 + a^2)(x^2 + b^2)}$ is-

(A) $\frac{1}{b^2 - a^2} \left[b \tan^{-1} \frac{x}{b} - a \tan^{-1} \frac{x}{a} \right] + c$

(B) $\frac{1}{b^2 - a^2} \left[a \tan^{-1} \frac{x}{b} - b \tan^{-1} \frac{x}{a} \right] + c$

(C) $\frac{1}{b^2 - a^2} \left[b \tan^{-1} \frac{x}{b} + a \tan^{-1} \frac{x}{a} \right] + c$

(D) None of these

Sol. Putting $x^2 = y$ in integrand, we obtain

$$\frac{y}{(y+a^2)(y+b^2)} = \frac{1}{b^2-a^2} \left[\frac{b^2}{y+b^2} - \frac{a^2}{y+a^2} \right]$$

$$\therefore I = \frac{1}{b^2-a^2} \cdot \left[\int \frac{b^2}{x^2+b^2} dx - \int \frac{a^2}{x^2+a^2} dx \right]$$

$$= \frac{1}{b^2-a^2} \left[b \tan^{-1} \frac{x}{b} - a \tan^{-1} \frac{x}{a} \right] + c$$

Ans.[A]

Ex.33 $\int \frac{dx}{3x^2 + 2x + 1}$ equals-

(A) $\frac{1}{\sqrt{2}} \tan^{-1} \left(\frac{3x+1}{\sqrt{2}} \right) + c$

(B) $\frac{1}{\sqrt{2}} \sin^{-1} \left(\frac{3x+1}{\sqrt{2}} \right) + c$

(C) $\frac{1}{\sqrt{2}} \cot^{-1} \left(\frac{3x+1}{\sqrt{2}} \right) + c$

(D) None of these

Sol. $I = \frac{1}{3} \int \frac{dx}{x^2 + \frac{2}{3}x + \frac{1}{3}}$

$$= \frac{1}{3} \int \frac{dx}{\left(x + \frac{1}{3} \right)^2 + \frac{2}{9}}$$

$$= \frac{1}{3} \times \frac{3}{\sqrt{2}} \tan^{-1} \left(\frac{x + \frac{1}{3}}{\sqrt{2}/3} \right) c$$

$$= \frac{1}{\sqrt{2}} \tan^{-1} \left(\frac{3x+1}{\sqrt{2}} \right) + c$$

Ans.[A]

Ex.34 $\int \sqrt{1+x-2x^2} dx$ equals-

(A) $\frac{1}{8} (4x-1) \sqrt{1+x-2x^2} + \frac{9\sqrt{2}}{32} \sin^{-1} \left(\frac{4x-1}{3} \right) + c$

(B) $\frac{1}{8} (4x+1) \sqrt{1+x-2x^2} - \frac{9\sqrt{2}}{32} \sin^{-1} \left(\frac{4x-1}{3} \right) + c$

(C) $\frac{1}{8} (4x-1) \sqrt{1+x-2x^2} + \frac{9\sqrt{2}}{32} \cos^{-1} \left(\frac{4x-1}{3} \right) + c$

(D) None of these

Sol. $I = \sqrt{2} \int \sqrt{\frac{1}{2} - \left(x^2 - \frac{x}{2} \right)} dx$

$$= \sqrt{2} \int \sqrt{\frac{9}{16} - \left(x - \frac{1}{4} \right)^2} dx$$

$$= \sqrt{2} \left[\frac{1}{2} \left(x - \frac{1}{4} \right) \sqrt{\frac{9}{16} - \left(x - \frac{1}{4} \right)^2} \right]$$

$$+ \frac{9}{32} \sin^{-1} \left\{ \frac{4}{3} \left(x - \frac{1}{4} \right) \right\} + c$$

$$= \frac{1}{8} (4x-1) \sqrt{1+x-2x^2} + \frac{9\sqrt{2}}{32} \sin^{-1} \left(\frac{4x-1}{3} \right) + c$$

Ans. [A]

Ex.35 $\int \frac{dx}{\sqrt{3-5x-x^2}}$ equals-

(A) $\sin^{-1} \left(\frac{2x+5}{\sqrt{37}} \right) + c$

(B) $\cos^{-1} \left(\frac{2x+5}{\sqrt{37}} \right) + c$

(C) $\sin^{-1} (2x+5) + c$

(D) None of these



Sol. $I = \int \frac{dx}{\sqrt{\frac{37}{4} - \left(x + \frac{5}{2}\right)^2}}$

$= \sin^{-1} \left(\frac{x + 5/2}{\sqrt{37}/2} \right) + C$

$= \sin^{-1} \left(\frac{2x + 5}{\sqrt{37}} \right) + C$

[A]

Ex.36 $\int \sqrt{e^{2x} - 1} dx$ is equal to-

(A) $\sqrt{e^{2x} - 1} + \sec^{-1} e^{2x} + C$

(B) $\sqrt{e^{2x} - 1} - \sec^{-1} e^{2x} + C$

(C) $\sqrt{e^{2x} - 1} - \sec^{-1} e^x + C$

(D) None of these

Sol. $\int \frac{e^{2x} - 1}{\sqrt{e^{2x} - 1}} dx$

$= \frac{1}{2} \int \frac{2e^{2x}}{\sqrt{e^{2x} - 1}} dx - \int \frac{e^x}{e^x \sqrt{e^{2x} - 1}} dx$

$= \sqrt{e^{2x} - 1} - \sec^{-1} e^x + C$

Ans.[C]

Ex.37 $\int \sqrt{\frac{e^x + a}{e^x - a}} dx$ is equal to-

(A) $\cosh^{-1} \left(\frac{e^x}{a} \right) + \sec^{-1} \left(\frac{e^x}{a} \right) + C$

(B) $\sinh^{-1} \left(\frac{e^x}{a} \right) + \sec^{-1} \left(\frac{e^x}{a} \right) + C$

(C) $\tanh^{-1} \left(\frac{e^x}{a} \right) + \cos^{-1} \left(\frac{e^x}{a} \right) + C$

(D) None of these

Sol. $\int \frac{e^x + a}{\sqrt{e^{2x} - a^2}} dx$

$= \int \frac{e^x}{\sqrt{e^{2x} - a^2}} dx + a \int \frac{e^x}{e^x \sqrt{e^{2x} - a^2}} dx$

$= \cosh^{-1} \left(\frac{e^x}{a} \right) + \sec^{-1} \left(\frac{e^x}{a} \right) + C$

Ans.[A]

Ex.38 $\int \frac{dx}{4\sin^2 x + 4\sin x \cos x + 5\cos^2 x}$ is equal to-

(A) $\tan^{-1} \left(\tan x + \frac{1}{2} \right) + C$

(B) $\frac{1}{4} \tan^{-1} \left(\tan x + \frac{1}{2} \right) + C$

(C) $4 \tan^{-1} \left(\tan x + \frac{1}{2} \right) + C$

(D) None of these

Sol. After dividing by $\cos^2 x$ to numerator and denominator of integration

$I = \int \frac{\sec^2 x dx}{4\tan^2 x + 4\tan x + 5}$

$= \int \frac{\sec^2 x dx}{(2\tan x + 1)^2 + 4}$

$= \frac{1}{2} \tan^{-1} \left(\frac{2\tan x + 1}{2} \right) + C$

Ans. [B]

Ex.39 $\int \left(\frac{1-x}{1+x} \right)^2 dx$ is equal to-

(A) $x - 4 \log(x+1) + \frac{4}{x+1} + C$

(B) $x - \log(x+1) + \frac{4}{x+1} + C$

(C) $x - 4 \log(x+1) - \frac{4}{x+1} + C$

(D) $x + \log(x+1) - \frac{4}{x+1} + C$

Sol. $\int \frac{[2-(x+1)]^2}{(x+1)^2} dx$

$= \int \left[\frac{4}{(x+1)^2} - \frac{4}{x+1} + 1 \right] dx$

$= -\frac{4}{x+1} - 4 \log(x+1) + x + C$

Ans.

[C]

Ex.40 $\int \frac{e^x}{e^{2x} + 5e^x + 6}$ equals-



(A) $\log \left(\frac{e^x + 3}{e^x + 2} \right) + c$

(B) $\log \left(\frac{e^x + 2}{e^x + 3} \right) + c$

(C) $\frac{1}{2} \log \left(\frac{e^x + 2}{e^x + 3} \right) + c$

(D) None of these

Sol. Put $e^x = t \Rightarrow e^x dx = dt$

$$\therefore I = \int \frac{dt}{t^2 + 5t + 6} = \int \frac{dt}{(t+2)(t+3)}$$

$$= \int \left(\frac{1}{t+2} - \frac{1}{t+3} \right) dt$$

$$= \log \left(\frac{t+2}{t+3} \right) + c = \log \left(\frac{e^x + 2}{e^x + 3} \right) + c$$

Ans. [B]

Ex.41 $\int \frac{dx}{x + \sqrt{x}}$ equals-

- (A) $2\log(\sqrt{x} - 1) + c$ (B) $2\log(\sqrt{x} + 1) + c$
 (C) $\tan^{-1} x + c$ (D) None of these

Sol. $I = \int \frac{dx}{x + \sqrt{x}}$

$$= \int \frac{2t dt}{t^2 + t} \text{ where } t^2 = x$$

$$= 2 \int \frac{dt}{t+1} = 2 \log(\sqrt{x} + 1) + c$$

Ans. [B]

Ex.42 $I = \int \frac{4e^x + 6e^{-x}}{9e^x - 4e^{-x}} dx$ is equal to-

- (A) $\frac{19}{36}x + \frac{35}{36}\log(9e^x - 4e^{-x}) + c$
 (B) $-\frac{19}{36}x + \frac{35}{36}\log(9e^x - 4e^{-x}) + c$
 (C) $\frac{1}{36}x + \frac{1}{36}\log(9e^x - 4e^{-x}) + c$
 (D) None of these

Sol. Suppose $4e^x + 6e^{-x} = A(9e^x - 4e^{-x}) + B(9e^x + 4e^{-x})$

By comparing $4 = 9A + 9B$,

$$6 = -4A + 4B$$

$$\text{or } A + B = \frac{4}{9}, -A + B = \frac{3}{2}$$

$$\text{After solving } A = -\frac{19}{36}, B = \frac{35}{36}$$

$$\therefore I = \int \left[-\frac{19}{36} + \frac{35}{36} \left(\frac{9e^x + 4e^{-x}}{9e^x - 4e^{-x}} \right) \right] dx$$

$$= -\frac{19}{36}x + \frac{35}{36}\log(9e^x - 4e^{-x}) + c$$

Ans. [B]

Ex.43 $\int \frac{\sin^{-1} \sqrt{x}}{\sqrt{1-x}} dx$ equals-

- (A) $2[\sqrt{x} - \sqrt{1-x} \sin^{-1} \sqrt{x}] + c$
 (B) $2[\sqrt{x} + \sqrt{1-x} \sin^{-1} \sqrt{x}] + c$
 (C) $[\sqrt{x} - \sqrt{1-x} \sin^{-1} \sqrt{x}] + c$
 (D) None of these

Sol. Let $x = \sin^2 t$, then

$$dx = 2 \sin t \cos t dt$$

$$\begin{aligned} \therefore I &= \int \frac{t}{\cos t} \cdot 2 \sin t \cos t dt \\ &= 2 \int t \sin t dt \\ &= 2[-t \cos t + \sin t] + c \\ &= 2[\sqrt{x} - \sqrt{1-x} \sin^{-1} \sqrt{x}] + c \end{aligned}$$

Ans. [A]

Ex.44 $\int \frac{\sqrt{x} - \sqrt{a}}{\sqrt{x+a}} dx$ equals-

- (A) $\sqrt{x^2 + ax} - 2\sqrt{ax + a^2} - a \cosh^{-1} \left(\sqrt{\frac{x+a}{a}} \right) + c$
 (B) $\sqrt{x^2 + ax} + \sqrt{ax + a^2} - a \cosh^{-1} \left(\sqrt{\frac{x+a}{a}} \right) c$
 (C) $\sqrt{x^2 + ax} - 2\sqrt{ax + a^2} + a \cosh^{-1} \left(\sqrt{\frac{x+a}{a}} \right) + c$
 (D) None of these

Sol. Let $x = a \tan^2 \theta \Rightarrow dx = 2a \tan \theta \sec^2 \theta d\theta$

$$\therefore I = \int \frac{\sqrt{a}(\tan \theta - 1) \cdot 2a \tan \theta \sec^2 \theta}{\sqrt{a} \sec \theta} d\theta$$

$$= 2a \left[\int \tan^2 \theta \sec \theta d\theta - \int \sec \theta \tan \theta d\theta \right]$$

$$\begin{aligned}
&= 2a \left[\int \sqrt{\sec^2 \theta - 1} \tan \theta \sec \theta d\theta - \sec \theta \right] \\
&= 2a \int \sqrt{t^2 - 1} dt - 2a \sec \theta + c \quad [\text{Where } \sec \theta = t] \\
&= 2a \left[\frac{t}{2} \sqrt{t^2 - 1} - \frac{1}{2} \cosh^{-1}(t) \right] - 2a \sqrt{\frac{a+x}{a}} + c \\
&= a \sqrt{\frac{x+a}{a}} \cdot \frac{x}{a} - a \cosh^{-1} \left(\sqrt{\frac{x+a}{a}} \right) \\
&\quad - 2 \sqrt{ax + a^2} + c
\end{aligned}$$

$$= \sqrt{x^2 + ax} - 2 \sqrt{ax + a^2} - a \cosh^{-1} \left(\sqrt{\frac{x+a}{a}} \right) + c$$

Ans. [A]

Ex.45 $\int \frac{x^5}{\sqrt{1+x^3}} dx$ equals-

(A) $\frac{2}{9}(x^3 - 2)\sqrt{1+x^3} + c$

(B) $\frac{2}{9}(x^3 + 2)\sqrt{1+x^3} + c$

(C) $(x^3 + 2)\sqrt{1+x^3} + c$

(D) None of these

Sol. Put $1+x^3 = t^2 \Rightarrow 3x^2 dx = 2t dt$

$$\therefore I = \int \frac{x^3}{\sqrt{1+x^3}} (x^2 dx) = \frac{2}{3} \int (t^2 - 1) dt$$

$$= \frac{2}{3} \left[\frac{t^3}{3} - t \right] + c$$

$$= \frac{2}{3} \left[\frac{1}{3}(1+x^3)^{3/2} - \sqrt{1+x^3} \right] + c$$

$$= \frac{2}{9} \sqrt{1+x^3} (1+x^3 - 3) + c$$

$$= \frac{2}{9} (x^3 - 2) \sqrt{1+x^3} + c$$

Ans.

[A]

Ex. 46 $\int \frac{e^{2\tan^{-1}x}(1+x)^2}{(1+x^2)} dx$ is equal to-

(A) $x e^{2\tan^{-1}x} + c$ (B) $x e^{2\tan^{-1}x} + c$

(C) $2x e^{2\tan^{-1}x} + c$ (D) None of these

Sol. Putting the value of $2 \tan^{-1} x = t$

$$\begin{aligned}
I &= \frac{1}{2} \int e^t \{1 + \tan(t/2)\}^2 dt \\
&= \frac{1}{2} \int e^t \left[\sec^2 \frac{t}{2} + 2 \tan \frac{t}{2} \right] dt \\
&= \frac{1}{2} e^t (2 \tan t/2) \\
&= e^t \tan \frac{t}{2} = x e^{2\tan^{-1}x} + c
\end{aligned}$$

Ans.

[B]

Ex.47 If $I = \int \cos^{-1} \sqrt{x} dx$ and

$$J = \int \frac{\sin^{-1} \sqrt{x} - \cos^{-1} \sqrt{x}}{\sin^{-1} \sqrt{x} + \cos^{-1} \sqrt{x}} dx, \text{ then } J \text{ equals-}$$

(A) $x - 4I$ (B) $x + I$

(C) $x - \frac{4}{\pi} I$ (D) $\frac{\pi}{4}$

Sol. Here

$$J = \frac{2}{\pi} \int \{\sin^{-1} \sqrt{x} - \cos^{-1} \sqrt{x}\} dx$$

$$= \frac{2}{\pi} \left(\frac{\pi}{2} - 2 \cos^{-1} \sqrt{x} \right) dx$$

$$[\because \sin^{-1} \sqrt{x} + \cos^{-1} \sqrt{x} = \frac{\pi}{2}]$$

$$= \int dx - \frac{4}{\pi} \int \cos^{-1} \sqrt{x} dx$$

$$= x - \frac{4}{\pi} I.$$

Ans. [C]

Ex.48 Which value of the constant of integration will make the integral of $\sin 3x \cos 5x$ zero at $x = 0$

(A) 0 (B) $-3/16$

(C) $-5/16$ (D) $1/8$

Sol. $I = \frac{1}{2} \int (\sin 8x - \sin 2x) dx$

$$= \frac{1}{2} \left[-\frac{\cos 8x}{8} + \frac{\cos 2x}{2} \right] + c$$

$$\text{At } x = 0, I = -\frac{1}{16} + \frac{1}{4} + c$$

$$\therefore I = 0 \Rightarrow c = -3/16$$

Ans. [B]

Ex.49 If $\int \frac{dx}{1+\sin x} = \tan \left(\frac{x}{2} + a \right) + c$, then value of a is



(A) $\frac{\pi}{4}$

(B) $-\frac{\pi}{4}$

(C) π

(D) $\frac{\pi}{2}$

Sol. $I = \int \frac{dx}{1 + \sin x}$

$$= \int \frac{dx}{1 + \cos\{(\pi/2) - x\}}$$

$$= \frac{1}{2} \int \sec^2\left(\frac{\pi}{4} - \frac{x}{2}\right) dx$$

$$= -\tan\left(\frac{\pi}{4} - \frac{x}{2}\right) + c = \tan\left(-\frac{\pi}{4} + \frac{x}{2}\right) + c$$

$$\therefore \alpha = -\frac{\pi}{4}$$

Ans. [B]

Ex. 50 If $\int \frac{2x+3}{(x-1)(x^2+1)} dx = \log[(x-1)^{5/2}(x^2+1)^a] - \frac{1}{2} \tan^{-1} x + k$ where k is any arbitrary constant, then a is equal to

(A) $5/4$

(B) $-5/3$

(C) $-5/6$

(D) $-5/4$

Sol.

$$\text{Let } I = \frac{2x+3}{(x-1)(x^2+1)} = \frac{A}{x-1} + \frac{Bx+C}{x^2+1}$$

$$\Rightarrow 2x+3 = A(x^2+1) + (Bx+C)(x-1) \dots (1)$$

Now putting $x = 1$, we get $5 = 2A \Rightarrow A = 5/2$

Equating coefficients of similar terms on both sides of (1),

we get,

$$-B + C = 2, A - C = 3$$

$$\Rightarrow C = 5/2 - 3 = -1/2$$

$$B = -1/2 - 2 = -5/2$$

$$\therefore I = \frac{5}{2} \int \frac{dx}{x-1} + \int \frac{-\frac{5}{2}x - \frac{1}{2}}{x^2+1} dx$$

$$= \frac{5}{2} \log(x-1) - \frac{5}{4} \int \frac{2x}{x^2+1} dx - \frac{1}{2} \int \frac{1}{x^2+1} dx$$

$$= \frac{5}{2} \log(x-1) - \frac{5}{4} \log(x^2+1) - \frac{1}{2} \tan^{-1} x + c$$

$$= \log [(x-1)^{5/2} (x^2+1)^{-5/4}] - \frac{1}{2} \tan^{-1} x + c$$

$$\therefore a = -5/4.$$

Ans. [D]