

SOLVED EXAMPLES

Ex.1 $\int_0^1 \frac{6x^2 + 1}{4x^3 + 2x + 3} dx$ is equal to-

- (A) $-\frac{1}{2} \log 3$ (B) $\frac{1}{2} \log 3$

- (C) $2 \log 3$ (D) None of these

Sol. Let $4x^3 + 2x + 3 = t$ $\therefore 2(6x^2 + 1)dx = dt$
Limits - at $x = 0; t = 3$, at $x = 1; t = 9$

$$\begin{aligned}\therefore I &= \int_3^9 \frac{1}{2} \frac{dt}{t} = \frac{1}{2} [\log t]_3^9 \\ &= \frac{1}{2} [\log 9 - \log 3] = \frac{1}{2} \log 3\end{aligned}$$

Ans. [B]

Ex.2 $\int_0^1 \frac{x}{1+x^4} dx$ is equal to -

- (A) $\frac{\pi}{2}$ (B) $\frac{\pi}{4}$ (C) $\frac{\pi}{8}$ (D) π

$$\begin{aligned}I &= \frac{1}{2} \int_0^1 \frac{2x}{1+(x^2)^2} dx \\ &= \frac{1}{2} [\tan^{-1} x^2]_0^1 \\ &= \frac{1}{2} [\tan^{-1} 1 - \tan^{-1} 0] \\ &= \frac{1}{2} \left[\frac{\pi}{4} - 0 \right] = \frac{\pi}{8}\end{aligned}$$

Ans. [C]

Ex.3 $\int_2^4 \frac{\sqrt{x^2 - 4}}{x} dx$ is equal to

- (A) $2(3\sqrt{3} - \pi)$ (B) $2\sqrt{3} - \pi$
(C) $\frac{2}{3}(3\sqrt{3} - \pi)$ (D) π

Sol. Put $x = 2 \sec t$, then

$$\begin{aligned}I &= \int_0^{\pi/3} \frac{2 \tan t}{2 \sec t} \cdot 2 \sec t \tan t dt \\ &= 2 \int_0^{\pi/3} \tan^2 t dt \\ &= 2 \int_0^{\pi/3} (\sec^2 t - 1) dt = 2[\tan t - t]_0^{\pi/3} \\ &= 2[\sqrt{3} - \pi/3] = \frac{2}{3}(3\sqrt{3} - \pi)\end{aligned}$$

Ans. [C]

Ex.4 $\int_0^{\pi/4} \frac{\sin \sqrt{x}}{\sqrt{x}} dx$ is equal to

- (A) 2 (B) 1
(C) $\pi/4$ (D) $\pi^2/8$

Sol. $\sqrt{x} = t, \frac{1}{\sqrt{x}} dx = dt$

$$\therefore I = 2 \int_0^{\pi/2} \sin t dt = 2(-\cos t)_0^{\pi/2} = 2(0 + 1) = 2$$

Ans. [A]

Ex.5 If $f(x) = \begin{cases} 2x+1, & 0 < x < 1 \\ x^2+2, & 1 \leq x < 2 \end{cases}$, then the value of

$$\int_0^2 f(x) dx$$

- (A) $-\frac{19}{3}$ (B) $\frac{19}{3}$

- (C) $\frac{3}{19}$ (D) None of these

$$\begin{aligned}\text{Sol. } \int_0^2 f(x) dx &= \int_0^1 f(x) dx + \int_1^2 f(x) dx \\ &= \int_0^1 (2x+1) dx + \int_1^2 (x^2+2) dx \\ &= \left[x^2 + x \right]_0^1 + \left[\frac{x^3}{3} + 2x \right]_1^2 \\ &= 2 - 0 + \left(\frac{20}{3} - \frac{7}{3} \right) = \frac{19}{3}\end{aligned}$$

Ans. [B]

Ex.6 $\int_0^{\pi/2} \log \sin x dx$ is equal to-

- (A) $\frac{\pi}{2} \log 2$ (B) $-\frac{\pi}{2} \log 2$

- (C) $\frac{\pi}{2} \log_{10} 2$ (D) $-\frac{\pi}{2} \log_{10} 2$

Sol. $I = \int_0^{\pi/2} \log \sin x dx$... (1)

$I = \int_0^{\pi/2} \log \cos x dx$ (by p-4) ... (2)

$$\therefore 2I = \int_0^{\pi/2} \log(\sin x \cos x) dx$$

$$\begin{aligned}
&= \int_0^{\pi/2} \log\left(\frac{\sin 2x}{2}\right) dx \\
&= \int_0^{\pi/2} \log \sin 2x dx - \frac{\pi}{2} \log 2 \\
&= \frac{1}{2} \int_0^{\pi} \log \sin t dt - \frac{\pi}{2} \log 2, \\
&\quad \text{where } t = 2x \\
&= 2 \frac{1}{2} \int_0^{\pi/2} \log \sin t dt - \frac{\pi}{2} \log 2 = 1 - \frac{\pi}{2} \log 2
\end{aligned}$$

$$\Rightarrow I = -\frac{\pi}{2} \log 2 \quad \text{Ans.[B]}$$

- Ex.7** $\int_0^{\pi/2} \frac{\sqrt{\sin x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx$ is equal to
 (A) $\pi/2$ (B) $\pi/4$
 (C) π (D) 2π

Sol. Using prop. P-4, we have

$$I = \int_0^{\pi/2} \frac{\sqrt{\cos x}}{\sqrt{\cos x} + \sqrt{\sin x}} dx$$

Adding it to given integral we have

$$\begin{aligned}
2I &= \int_0^{\pi/2} dx = [x]_0^{\pi/2} = \pi/2 \\
\therefore I &= \pi/4 \quad \text{Ans.[B]}
\end{aligned}$$

- Ex.8** If $f(x)$ is an odd function of x , then

$$\begin{aligned}
&\int_{-\pi/2}^{\pi/2} f(\cos x) dx \text{ is equal to} \\
&\quad \text{(A) 0} \quad \text{(B) } \int_0^{\pi/2} f(\cos x) dx \\
&\quad \text{(C) } 2 \int_0^{\pi/2} f(\sin x) dx \quad \text{(D) } \int_0^{\pi} f(\cos x) dx
\end{aligned}$$

Sol. Here $f(\cos x)$ will be even function of x ,

$$\begin{aligned}
I &= \int_{-\pi/2}^{\pi/2} f(\cos x) dx = 2 \int_0^{\pi/2} f(\cos x) dx \\
&= 2 \int_0^{\pi/2} f(\sin x) dx \quad \text{Ans.[C]}
\end{aligned}$$

- Ex.9** The value of the integral

$$\int_{-4}^4 (ax^3 + bx + c) dx \text{ depend on}$$

- (A) b and c (B) a, b and c
 (C) only c (D) a and c

$$\begin{aligned}
\text{Sol.} \quad I &= \int_{-4}^4 (ax^3 + bx) dx + \int_{-4}^4 c dx \\
&= 0 + 2 \int_0^4 c dx \quad (\text{by P-5}) \\
&= 2c[x]_0^4 = 8c
\end{aligned}$$

Hence the value of I depends on c.

Ans.[C]

- Ex.10** If $f(x) = \frac{x \cos x}{1 + \sin^2 x}$, then $\int_{-\pi}^{\pi} f(x) dx$ equals-

- (A) $\pi/4$ (B) $\pi/2$
 (C) π (D) 0

$$\text{Sol.} \quad \text{Since } f(-x) = \frac{-x \cos(-x)}{1 + \sin^2(\pi - x)}$$

$$= \frac{-x \cos x}{1 + \sin^2 x} = -f(x)$$

$$\therefore I = \int_{-\pi}^{\pi} f(x) dx = 0 \quad \text{Ans.[D]}$$

- Ex.11** $\int_0^{\pi/2} \sin^2 x \cos^3 x dx$ equals-

- (A) 1 (B) $2/5$
 (C) $2/15$ (D) $4/15$

Sol. Using Walli's formula, we get

$$I = \frac{1.2}{5.3.1} = \frac{2}{15} \quad \text{Ans.[C]}$$

- Ex.12** $\int_{\pi/4}^{3\pi/4} \frac{\phi}{1 + \sin \phi} d\phi$ equals-

- (A) $\pi(\sqrt{2} - 1)$ (B) $\pi(\sqrt{2} + 1)$
 (C) $\pi(2 - \sqrt{2})$ (D) None of these

$$\text{Sol.} \quad I = \int_{\pi/4}^{3\pi/4} \frac{\phi}{1 + \sin \phi} d\phi \quad \dots(1)$$

$$\Rightarrow I = \int_{\pi/4}^{3\pi/4} \frac{\pi - \phi}{1 + \sin(\pi - \phi)} d\phi \quad (\text{by P-8})$$

$$I = I_1 + I_2$$

Putting $x = \frac{1}{t}$ in second integrand

$$dx = -\frac{1}{t^2} dt$$

$$\therefore I_2 = \int_1^0 \frac{\frac{1}{t} \log\left(\frac{1}{t}\right)}{\left(1 + \frac{1}{t^2}\right)^2} \left(-\frac{1}{t^2}\right) dt$$

$$= - \int_0^1 \frac{t \log t}{(1+t^2)^2} dt = -I_1$$

$$I = I_2 + I_1 = -I_1 + I_1 = 0$$

Ans.[A]

Ex.20 $\int_0^\pi x \sin^4 x dx$ is equal to-

- (A) $3\pi/16$ (B) $3\pi^2/16$
 (C) $16\pi/3$ (D) $16\pi^2/3$

Sol. $I = \int_0^\pi x \sin^4 x dx$... (1)

$$I = \int_0^\pi (\pi - x) \sin^4(\pi - x) dx$$

$$I = \int_0^\pi (\pi - x) \sin^4 x dx$$

$$\therefore 2I = \pi \int_0^\pi \sin^4 x dx$$

$$\Rightarrow 2I = \pi \int_0^\pi \sin^4 x dx \quad [\text{from property P-6}]$$

$$\Rightarrow I = \pi \cdot \frac{3.1}{4.2} \cdot \frac{\pi}{2} = \frac{3\pi^2}{16}$$

Ans.[B]

Ex.21 $\int_1^2 \log x dx$ equals-

- (A) $2 \log 2$ (B) $\log\left(\frac{2}{e}\right)$
 (C) $\log\left(\frac{4}{e}\right)$ (D) None of these

Sol. $I = \int_1^2 1 \cdot \log x dx$ equals

(Integrating by parts by taking 1 as a second function)

$$= \{x \cdot \log x\}_1^2 - \int_1^2 \left(\frac{1}{x} \cdot x \right) dx$$

$$= (2 \log 2 - 1 \log 1) - [x]_1^2$$

$$= (2 \log 2 - 0) - (2 - 1)$$

$$= \log 4 - \log e = \log\left(\frac{4}{e}\right)$$

Ans.[C]

Ex.22 $\int_0^{\pi/2} \frac{2^{\sin x}}{2^{\sin x} + 2^{\cos x}} dx$ equals-

- (A) 2 (B) π

- (C) $\frac{\pi}{4}$ (D) $\frac{\pi}{2}$

Sol. $I = \int_0^{\pi/2} \frac{2^{\sin x}}{2^{\sin x} + 2^{\cos x}} dx$

$$I = \int_0^{\pi/2} \frac{2^{\sin(\pi/2-x)}}{2^{\sin(\pi/2-x)} + 2^{\cos(\pi/2-x)}} dx$$

$$= \int \frac{2^{\cos x}}{2^{\cos x} + 2^{\sin x}} dx$$

$$2I = \int_0^{\pi/2} dx = \frac{\pi}{2} \Rightarrow I = \frac{\pi}{4}$$

Ans.[C]

Ex.23 $\int_0^x f(t) dt = x + \int_x^1 t f(t) dt$ then $f(1)$ is equal to-

- (A) $\frac{1}{2}$ (B) 0

- (C) 1 (D) $-\frac{1}{2}$

Sol. $\int_0^x f(t) dt = x + \int_x^1 t f(t) dt$

$$\Rightarrow f(x) = 1 + (0 - xf(x)) \quad [\text{diff. w.r.t. } x]$$

$$\Rightarrow f(x) = 1 - xf(x)$$

$$\Rightarrow f(1) = 1 - 1 \cdot f(1)$$

$$\Rightarrow f(1) = \frac{1}{2}$$

Ans.[A]

Ex.24 If $f(3-x) = f(x)$ then $\int_1^2 xf(x) dx$ equals-

- (A) $\frac{3}{2} \int_1^2 f(2-x) dx$ (B) $\frac{3}{2} \int_1^2 f(x) dx$
 (C) $\frac{1}{2} \int_1^2 f(x) dx$ (D) None of these

Sol. Let $x = 3 - y$

$$\begin{aligned} I &= \int_2^1 (3-y)f(3-y)(-dy) \\ &= \int_1^2 (3-x)f(3-x)dx \\ &= \int_1^2 (3-x)f(x)dx \quad [\because f(3-x) = f(x)] \\ &= 3 \int_1^2 f(x)dx - I \\ I &= \frac{3}{2} \int_1^2 f(x)dx \end{aligned}$$

Ans.[B]

Ex.25 $\int_0^1 \frac{x \sin^{-1} x}{\sqrt{1-x^2}} dx$ is equal to-

- (A) $\pi/2$ (B) $\pi/4$
 (C) 0 (D) 1

Sol. Put $\sin^{-1} x = t$, $\frac{dx}{\sqrt{1-x^2}} = dt$ then

$$\therefore I = \int_0^{\pi/2} t \sin t dt = [t(-\cos t)]_0^{\pi/2} + [\sin x]_0^{\pi/2} = 1$$

Ans.[C]

Ex.26 The value of the integral $\int_0^{\pi/4} \frac{\sin \theta + \cos \theta}{9+16 \sin 2\theta} d\theta$ is

- (A) $\log 3$ (B) $\log 2$
 (C) $\frac{1}{20} \log 3$ (D) $\frac{1}{20} \log 2$

Sol. Here

$$\begin{aligned} I &= \int_0^{\pi/4} \frac{\sin \theta + \cos \theta}{9+16(\sin 2\theta+1-1)} d\theta \\ &= \int_0^{\pi/4} \frac{\sin \theta + \cos \theta}{25-16(1-\sin 2\theta)} d\theta \\ &= \frac{1}{16} \int_0^{\pi/4} \frac{\sin \theta + \cos \theta}{(25/16)-(\sin \theta-\cos \theta)^2} d\theta \end{aligned}$$

$$= \frac{1}{16} \int_{-1}^0 \frac{dt}{(25/16)-t^2}, \text{ where } (\sin \theta - \cos \theta) = t$$

$$\begin{aligned} &= \frac{1}{16} \times \frac{1}{2 \times 5/4} \left[\log \frac{(5/4)+t}{(5/4)-t} \right]_{-1}^0 \\ &= \frac{1}{40} \left[\log 1 - \log \frac{1/4}{9/4} \right] = \frac{1}{20} \log 3 \end{aligned}$$

Ans.[C]

Ex.27 $\int_{-\pi/2}^{\pi/2} \sin^2 x \cos^2 x (\sin x + \cos x) dx$ is equal to-

- (A) 2/15 (B) 4/15
 (C) 2/5 (D) 8/15

Sol. $I = \int_{-\pi/2}^{\pi/2} \sin^3 x \cos^2 x dx + \int_{-\pi/2}^{\pi/2} \sin^2 x \cos^3 x dx$
 (by P-5)

$$\begin{aligned} &= 0 + 2 \int_0^{\pi/2} \sin^2 x \cos^3 x dx \\ &= 2 \cdot \frac{1.2}{5.3.1} = \frac{4}{15} \end{aligned}$$

Ans.[B]

Ex.28 $\int_0^{2a} \frac{f(x)}{f(x)+f(2a-x)} dx$ is equal to-

- (A) a (B) -a
 (C) 0 (D) None of these

Sol. Using P-4, given integral becomes

$$I = \int_0^{2a} \frac{f(2a-x)}{f(2a-x)+f(x)} dx$$

Adding it with the given integral, we get

$$2I = \int_0^{2a} 1 dx = 2a \Rightarrow I = a \quad \text{Ans.[A]}$$

Ex.29 $\int_{-1}^{3/2} |x \sin \pi x| dx$ is equal to

- (A) $\frac{4}{\pi}$ (B) $\frac{3}{\pi} + \frac{1}{\pi^2}$
 (C) $\frac{3}{\pi^2} + \frac{1}{\pi}$ (D) None of these

Sol. Obviously

$$|x \sin \pi x| = \begin{cases} x \sin \pi x, & -1 < x < 1 \\ -x \sin \pi x, & 1 < x < 3/2 \end{cases}$$

$$\begin{aligned}
I &= \int_{-1}^1 x \sin \pi x dx + \int_1^{3/2} (-x \sin \pi x) dx \\
&= 2 \int_0^{1/2} x \sin \pi x dx - \int_1^{3/2} x \sin \pi x dx \\
&= 2 \left[-\frac{x}{\pi} \cos \pi x + \frac{1}{\pi^2} \sin \pi x \right]_0^1 \\
&\quad - \left[-\frac{x}{\pi} \cos \pi x + \frac{1}{\pi^2} \sin \pi x \right]_1^{3/2} \\
&= 2 \left(-\frac{\cos \pi}{\pi} \right) - \left(\frac{\sin(3\pi/2)}{\pi^2} + \frac{\cos \pi}{\pi} \right) \\
&= \frac{3}{\pi} + \frac{1}{\pi^2} \tag{Ans.[B]}
\end{aligned}$$

Ex.30 The value of $\int_0^{100\pi} \sqrt{1-\cos 2x} dx$ is
 (A) $100\sqrt{2}$ (B) $200\sqrt{2}$
 (C) $50\sqrt{2}$ (D) 0

Sol. $I = \sqrt{2} \int_0^{100\pi} |\sin x| dx$
 $= 100\sqrt{2} \int_0^{\pi} |\sin x| dx$
 $= 100\sqrt{2} \int_0^{\pi} \sin x dx = 100\sqrt{2} [-\cos x]_0^{\pi}$
 $= 200\sqrt{2} \tag{Ans.[B]}$

Ex.31 $\int_0^{2\pi} \frac{x \sin^{2n} x}{\sin^{2n} x + \cos^{2n} x} dx$ ($n \in \mathbb{N}$) is equal to-
 (A) π^2 (B) $2\pi^2$
 (C) π (D) 2π

Sol. $I = \int_0^{2\pi} \frac{x \sin^{2n} x}{\sin^{2n} x + \cos^{2n} x} dx$
 $= \int_0^{2\pi} \frac{(2\pi-x) \sin^{2n} (2\pi-x)}{\sin^{2n} x (2\pi-x) + \cos^{2n} (2\pi-x)} dx$
 (By P-4)

$$\begin{aligned}
&= \int_0^{2\pi} \frac{(2\pi-x) \sin^{2n} x}{\sin^{2n} x + \cos^{2n} x} dx \\
\therefore 2I &= 2\pi \int_0^{2\pi} \frac{\sin^{2n} x}{\sin^{2n} x + \cos^{2n} x} dx
\end{aligned}$$

$$\begin{aligned}
\Rightarrow I &= 4\pi \int_0^{\pi/2} \frac{\sin^{2n} x}{\sin^{2n} x + \cos^{2n} x} dx \\
&= 4\pi(\pi/4) = \pi^2. \tag{Ans.[A]}
\end{aligned}$$

Ex.32 $\int_0^{\pi/2} \frac{dx}{1+2\sin x+\cos x}$ equals-
 (A) $(1/2) \log 3$ (B) $\log 3$
 (C) $(4/3) \log 3$ (D) None of these

Sol. Here

$$\begin{aligned}
I &= \int_0^{\pi/2} \frac{dx}{1+2 \frac{2\tan(x/2)}{1+\tan^2(x/2)} + \frac{1-\tan^2(x/2)}{1+\tan^2(x/2)}} \\
&= \int_0^{\pi/2} \frac{\sec^2(x/2)}{2\{1+2\tan(x/2)\}} dx \\
\text{Let } 1+2\tan(x/2) &= t, \text{ then} \\
\sec^2(x/2) dx &= dt \\
\therefore I &= \frac{1}{2} \int_1^3 \frac{dt}{t} = \frac{1}{2} (\log t)_1^3 \\
&= \frac{1}{2} \log 3 \tag{Ans.[A]}
\end{aligned}$$

Ex.33 $\int_0^{\pi/2} \frac{\sin 2x}{a \cos^2 x + b \sin^2 x} dx$ -
 (A) $\frac{1}{b-a} \log \left(\frac{b}{a} \right)$ (B) $\frac{1}{b+a} \log \left(\frac{b}{a} \right)$
 (C) $\frac{1}{b-a} \log \left(\frac{a}{b} \right)$ (D) $\frac{1}{b+a} \log \left(\frac{a}{b} \right)$

Sol. $I = \left(\frac{1}{b-a} \right) \int_0^{\pi/2} \frac{(b-a) 2 \sin x \cos x}{a \cos^2 x + b \sin^2 x} dx$
 $= \frac{1}{b-a} \left[\log(a \cos^2 x + b \sin^2 x) \right]_0^{\pi/2}$
 $= \frac{1}{(b-a)} (\log b - \log a)$
 $= \frac{1}{b-a} \log \left(\frac{b}{a} \right) \tag{Ans.[A]}$

Ex.34 $\int_0^{\pi/2} (2 \log \sin x - \log \sin 2x) dx$ equals-
 (A) $\pi \log 2$ (B) $-\pi \log 2$

(C) $(\pi/2) \log 2$ (D) $-(\pi/2) \log 2$

Sol. $I = \int_0^{\pi/2} (2 \log \sin x - \log 2 \sin x \cos x) dx$

$$= \int_0^{\pi/2} (2 \log \sin x - \log 2 - \log \sin x - \log \cos x) dx$$

$$= \int_0^{\pi/2} \log \sin x dx - \int_0^{\pi/2} \log 2 dx - \int_0^{\pi/2} \log \cos x dx$$

$$= -(\pi/2) \log 2.$$

Ans.[D]

Ex.35 $\int_0^{\pi/2} \frac{x \sin x \cos x}{\sin^4 x + \cos^4 x} dx$ equals-

(A) $\frac{\pi^2}{8}$ (B) $\frac{\pi^2}{16}$
 (C) $\frac{\pi^2}{4}$ (D) $\frac{\pi^2}{2}$

Sol. $I = \frac{\pi}{4} \int_0^{\pi/2} \frac{\sin x \cos x}{\sin^4 x + \cos^4 x} dx$

$$= \frac{\pi}{8} \int_0^{\pi/2} \frac{2 \sin x \cos x}{(\sin^2 x)^2 + (1 - \sin^2 x)^2} dx$$

Assume $\sin^2 x = t$
 $\therefore 2 \sin x \cos x dx = dt$

$$\therefore I = \frac{\pi}{8} \int \frac{dt}{t^2 + (1-t)^2}$$

$$I = \frac{\pi}{8} \int \frac{dt}{2t^2 - 2t + 1}$$

$$= \frac{\pi}{16} \int \frac{dt}{\left(t - \frac{1}{2}\right)^2 + \frac{1}{4}}$$

$$= \frac{\pi}{16} \cdot \frac{1}{\left(\frac{1}{2}\right)} \tan^{-1} \left[\frac{\left(t - \frac{1}{2}\right)}{\frac{1}{2}} \right]$$

$$= \frac{\pi}{8} [\tan^{-1}(2 \sin^2 x - 1)]_0^{\pi/2}$$

$$= \frac{\pi}{8} [\tan^{-1}(1) - \tan^{-1}(-1)]$$

$$= \frac{\pi}{8} \left[\frac{\pi}{4} - \left(-\frac{\pi}{4} \right) \right] = \frac{\pi^2}{16}$$

Ans.[B]

Ex. 36 $\int_0^{\pi/2} |\sin x - \cos x| dx$ equals-

(A) $2\sqrt{2}$ (B) $2(\sqrt{2} + 1)$
 (C) $2(\sqrt{2} - 1)$ (D) 0

Sol. $\because |\sin x - \cos x|$

$$= \begin{cases} -(\sin x - \cos x), & 0 < x < \pi/4 \\ (\sin x - \cos x), & \pi/4 < x < \pi/2 \end{cases}$$

$$\therefore I = \int_0^{\pi/4} -(\sin x - \cos x) dx + \int_{\pi/4}^{\pi/2} (\sin x - \cos x) dx$$

$$= [\cos x + \sin x]_0^{\pi/4} + [-\cos x - \sin x]_{\pi/4}^{\pi/2}$$

$$= \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} - 1 - 1 + \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}}$$

$$= 2\sqrt{2} - 2$$

Ans.[C]

Ex.37 The value of $\lim_{x \rightarrow 0} \frac{\int_0^x \cos t^2 dt}{x}$ is-

(A) 0 (B) 1
 (C) -1 (D) None of these

Sol. Let $f(x) = \int_0^x \cos t^2 dt$ and $g(x) = x$,
 $\therefore \lim_{x \rightarrow 0} \frac{f(x)}{g(x)} = \lim_{x \rightarrow 0} \frac{f'(x)}{g'(x)}$

$$\therefore \text{Given limit} = \lim_{x \rightarrow 0} \frac{\cos x^2 \cdot 1 - \cos 0 \cdot 0}{1}$$

$$\left[\text{since } \frac{d}{dx} \int_{\phi(x)}^{\psi(x)} f(t) dt = \int_{\phi(x)}^{\psi(x)} \frac{d}{dx} (f(t)) dt \right]$$

$$= f(\psi(x)\psi'(x)) - f(\phi(x)\phi'(x))$$

$\therefore \text{Given limit} = \cos 0 = 1.$

Ans.[B]

Ex.38 If $n \in \mathbb{Z}$, then

$$\int_0^{\pi} e^{\sin^2 x} \cos^3(2n+1)x dx -$$

(A) -1 (B) 0
 (C) 1 (D) π

Sol. Let $f(x) = e^{\sin^2 x} \cos^3(2n+1)x \, dx$

$$\begin{aligned} \Rightarrow f(\pi - x) &= e^{\sin^2(\pi-x)} \cos^3(2n+1)(\pi-x)dx \\ &= -e^{\sin^2 x} \cos^3(2n+1)x \\ &[\because (2n+1) \text{ is odd}] \\ &= -f(x) \end{aligned}$$

So by P-8, I = 0

Ans.[B]

Ex.39 The value of $\int_{-1/2}^{1/2} \left[\left(\frac{x+1}{x-1} \right)^2 + \left(\frac{x-1}{x+1} \right)^2 - 2 \right]^{1/2} dx$

equals-

- (A) $\log(4/3)$ (B) $2 \log(4/3)$
 (C) $4 \log(4/3)$ (D) $-4 \log(4/3)$

Sol. Here

$$\begin{aligned} I &= \int_{-1/2}^{1/2} \left[\left(\frac{x+1}{x-1} - \frac{x-1}{x+1} \right)^2 \right]^{1/2} dx \\ &= \int_{-1/2}^{1/2} \left| \frac{4x}{x^2-1} \right| dx = 2 \int_0^{1/2} \left| \frac{4x}{x^2-1} \right| dx \\ &= 8 \int_0^{1/2} \frac{x \, dx}{1-x^2} = -4 \left[\log(1-x^2) \right]_0^{1/2} \\ &= -4 \log\left(\frac{3}{4}\right) = 4 \log\left(\frac{4}{3}\right) \quad \text{Ans.[C]} \end{aligned}$$

Ex.40 $\int_0^1 \cot^{-1}(1-x+x^2) dx$ equals-

- (A) $\frac{\pi}{2} + \log 2$ (B) $\frac{\pi}{2} - \log 2$
 (C) $\pi - \log 2$ (D) None of these

Sol. $I = \int_0^1 \tan^{-1}\left(\frac{1}{1-x-x^2}\right) dx$

$$\begin{aligned} &= \int_0^1 \tan^{-1}\left(\frac{x+(1-x)}{1-x(1-x)}\right) dx \\ &= \int_0^1 [\tan^{-1} x + \tan^{-1}(1-x)] dx \\ &= \int_0^1 \tan^{-1} x \, dx + \int_0^1 \tan^{-1}(1-x) \, dx \end{aligned}$$

$$= 2 \int_0^1 \tan^{-1} x \, dx \quad [\text{By prov. IV}]$$

$$= 2 \left[x \tan^{-1} - \frac{1}{2} \log(1+x^2) \right]_0^1$$

$$= 2 \frac{\pi}{4} - \log 2 = \frac{\pi}{2} - \log 2 \quad \text{Ans.[B]}$$

Ex.41 $\int_0^{\pi/4} (\sqrt{\tan x} + \sqrt{\cot x}) dx$ is equal to-

- (A) $\pi/2$ (B) $\pi/\sqrt{2}$
 (C) $-\pi/2$ (D) $-\pi/\sqrt{2}$

Sol. Putting $\tan x = t^2$, then

$$\sec^2 x \, dx = 2t \, dt \Rightarrow dx = \frac{2t \, dt}{1+t^4}$$

$$\therefore I = \int_0^1 \left(t + \frac{1}{t} \right) \frac{2t \, dt}{1+t^4}$$

$$= 2 \int_0^1 \frac{t^2+1}{t^4+1} dt = 2 \int_0^1 \frac{1+1/t^2}{t^2+1/t^2} dt$$

$$= 2 \int_0^1 \frac{d(t-1/t)}{(t-1/t)^2+2}$$

$$= \sqrt{2} \left[\tan^{-1} \frac{1}{\sqrt{2}} \left(t - \frac{1}{t} \right) \right]_0^1$$

$$= \sqrt{2} [\tan^{-1} 0 - \tan^{-1} (-\infty)]$$

$$= \sqrt{2} (\pi/2) = \pi/\sqrt{2} \quad \text{Ans.[B]}$$

Ex.42 If $g(x) = \int_0^x \cos^4 t \, dt$, then $g(x+\pi)$ is equal to-

- (A) $g(x) + g(\pi)$ (B) $g(x) - g(\pi)$
 (C) $g(x)g(\pi)$ (D) $g(x)/g(\pi)$

Sol. $g(x+\pi) = \int_0^{x+\pi} \cos^4 t \, dt$

$$= \int_0^\pi \cos^4 t \, dt + \int_\pi^{x+\pi} \cos^4 t \, dt \quad [\text{by P-3}]$$

$$= \int_0^\pi \cos^4 t \, dt + I_2$$

Now in I_2 , put $t = \pi + \theta$, then

$$I_2 = \int_0^\pi \cos^4(\pi+\theta) d\theta$$

$$= \int_0^\pi \cos^4 \theta d\theta = \int_0^\pi \cos^4 t \, dt$$

$$\therefore g(x+\pi) = \int_0^\pi \cos^4 t \, dt + \int_0^x \cos^4 t \, dt$$

$$= g(x) + g(\pi) \quad \text{Ans.[A]}$$

Ex.43 $\int_{-\pi/2}^{\pi/2} \frac{\cos x}{1+e^x} dx$ is equal to-

- (A) 0 (B) 2
 (C) 1 (D) None of these

Sol. $I = \int_{-\pi/2}^0 \frac{\cos x}{1+e^x} dx + \int_0^{\pi/2} \frac{\cos x}{1+e^x} dx$

 $= - \int_{\pi/2}^0 \frac{\cos y}{1+e^{-y}} dy + \int_0^{\pi/2} \frac{\cos x}{1+e^x}$

(putting $x = -y$ in first integral)

 $= \int_0^{\pi/2} \frac{e^y \cos y}{1+e^y} dy + \int_0^{\pi/2} \frac{\cos x}{1+e^x} dx$
 $= \int_0^{\pi/2} \frac{e^x \cos x}{1+e^x} dx + \int_0^{\pi/2} \frac{\cos x}{1+e^x} dx$
 $= \int_0^{\pi/2} \frac{(e^x + 1)\cos x}{1+e^x} dx$
 $= \int_0^{\pi/2} \cos x dx = [\sin x]_0^{\pi/2} = 1$

Ans.[C]

Ex.44 $\int_{-1}^1 \frac{\sin x - x^2}{3 - |x|} dx$ is equal to-

$$\begin{aligned}
 \text{Sol. } I &= \int_{-1}^1 \frac{\sin x - x^2}{3 - |x|} dx \\
 &= \int_{-1}^1 \frac{\sin x}{3 - |x|} dx - \int_{-1}^1 \frac{x^2}{3 - |x|} dx \\
 &= 0 - 2 \int_0^1 \frac{x^2}{3 - |x|} dx
 \end{aligned}$$

[\because $\frac{\sin x}{3-|x|}$ is an odd and $\frac{x^2}{3-|x|}$ is an even]

function]

$$= -2 \int_0^1 \frac{x^2}{3-|x|} dx$$

Ex.45 $\int_{-4}^{-5} e^{(x+5)^2} dx + 3 \int_{1/3}^{2/3} e^{9(x-2/3)^2} dx$ is equal to-

- (A) e^5 (B) e^4
 (C) $3e^2$ (D) 0

Sol. Putting $x = -t - 4$ in first integral and

$$x = \frac{t}{3} + \frac{1}{3} \text{ in second integral}$$

$$I_1 = \int_{-4}^{-5} e^{(x+5)^2} dx = - \int_0^1 e^{(-t+1)^2} dt = - \int_0^1 e^{(t-1)^2} dt$$

$$\begin{aligned} I_2 &= 3 \int_{1/3}^{2/3} e^{9(x-2/3)^2} dx \\ &= 3 \int_0^1 e^{9(t/3-1/3)^2} dt = \int_0^1 e^{(t-1)^2} dt \end{aligned}$$

$$\therefore I = I_1 + I_2 = 0.$$

Ans.[D]

Ex.46 Let f be a positive function. If

$$I_1 = \int_{1-k}^k xf\{x(1-x)\}dx$$

$$I_2 = \int_{1-k}^k f[x(1-x)]dx$$

where $2k - 1 > 0$, then the value of I_1/I_2 is equal to-

Sol. Using property P – 8, we have

$$I_1 = \int_{1-k}^k (k+1-k-x) f[(k+1-k-x) \times (1-k-1+k+x)] dx$$

$$= \int_{1-k}^k (1-x) f[(1-x)(x)] dx$$

$$= \int_{1-k}^k f[x(1-x)]dx - \int_{1-k}^k xf[x(1-x)]dx$$

$$= I_2 - I_1$$

$$\Rightarrow 2I_1 = I_2 \therefore \frac{I_1}{I_2} = \frac{1}{2}$$

Ans.[C]

